

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 February 2002 (28.02.2002)

PCT

(10) International Publication Number  
**WO 02/16310 A1**

(51) International Patent Classification<sup>7</sup>: **C07C 233/15**,  
233/33, 233/48, 235/26, 237/14, A01N 37/18, 31/16,  
31/165

(74) Agent: **COHEN, Mark, S.**; Eitan, Pearl, Latzer & Cohen-  
Zedek, One Crystal Park, Suite 210, 2011 Crystal Drive,  
Arlington, VA 22202-3709 (US).

(21) International Application Number: PCT/US01/26328

(22) International Filing Date: 23 August 2001 (23.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/644,970 24 August 2000 (24.08.2000) US  
60/300,083 25 June 2001 (25.06.2001) US

(71) Applicant (*for all designated States except US*): **GTX, INC.** [US/US]; 3 North Dunlap, Van Fleet Building, Memphis, TN 38163 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **DALTON, James** [US/US]; 2706 Wellesley Drive, Columbus, OH 43221 (US). **MILLER, Duane, D.** [US/US]; 8706 Maple Creek Cove, Germantown, TN 38139 (US). **YIN, Donghua** [US/US]; 984 Chatham Lane, Apt B, Columbus, OH 43221 (US). **HE, Yali** [US/US]; 812-H South Parker Drive, Florence, SC 25901 (US).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: SELECTIVE ANDROGEN RECEPTOR MODULATORS AND METHODS OF USE THEREOF

(57) Abstract: This invention provides a novel class of androgen receptor targeting agents (ARTA). The agents define a new subclass of compounds which are tissue-selective androgen receptor modulators (SARM), which are useful for oral testosterone replacement therapy, male contraception, maintaining sexual desire in women, treating prostate cancer and imaging prostate cancer. These agents have an unexpected in-vivo activity for an androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor. These agents may be active alone or in combination with progestins or estrogens. The invention further provides a novel class of non-steroidal agonist compounds. The invention further provides compositions containing the selective androgen modulator compounds or the non-steroidal agonist compounds and methods of binding an androgen receptor, modulating spermatogenesis, treating and imaging prostate cancer, and providing hormonal therapy for androgen-dependent conditions.

WO 02/16310 A1

**SELECTIVE ANDROGEN RECEPTOR MODULATORS AND  
METHODS OF USE THEREOF**

**FIELD OF INVENTION**

5

[0001] The present invention relates to a novel class of tissue-selective androgen receptor targeting agents (ARTA) which demonstrate androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor. The agents define a new subclass of compounds which are tissue-selective androgen receptor modulators (SARM) which are  
10 useful for male hormone therapy such as oral testosterone replacement therapy, male contraception, maintaining sexual desire in women, treating prostate cancer, and imaging prostate cancer. These agents are also administered to a subject for the treatment of sarcopenia, lack of sexual libido, osteoporosis, erythropoiesis, and fertility. The agents may be used alone or in combination with a progestin or estrogen.

15

**BACKGROUND OF THE INVENTION**

[0002] The androgen receptor ("AR") is a ligand-activated transcriptional regulatory protein that mediates induction of male sexual development and function through its activity with endogenous androgens. Androgens are generally known as the male sex  
20 hormones. However, androgens also play a pivotal role in female physiology and reproduction. The androgenic hormones are steroids which are produced in the body by the testis and the cortex of the adrenal gland, or synthesized in the laboratory. Androgenic steroids play an important role in many physiologic processes, including the development and maintenance of male sexual characteristics such as muscle and bone  
25 mass, prostate growth, spermatogenesis, and the male hair pattern (Matsumoto, Endocrinol. Met. Clin. N. Am. 23:857-75 (1994). The endogenous steroidal androgens include testosterone and dihydrotestosterone ("DHT"). Testosterone is the principal steroid secreted by the testes and is the primary circulating androgen found in the plasma of males. Testosterone is converted to DHT by the enzyme 5 alpha-reductase in many  
30 peripheral tissues. DHT is thus thought to serve as the intracellular mediator for most androgen actions (Zhou, et al., Molec. Endocrinol. 9:208-18 (1995)). Other steroidal androgens include esters of testosterone, such as the cypionate, propionate, phenylpropionate, cyclopentylpropionate, isocarporate, enanthate, and decanoate esters, and other synthetic androgens such as 7-Methyl-Nortestosterone ("MENT") and its

acetate ester (Sundaram et al., "7 Alpha-Methyl-Nortestosterone(MENT): The Optimal Androgen For Male Contraception," Ann. Med., 25:199-205 (1993) ("Sundaram")). Because the AR is involved in male sexual development and function, the AR is a likely target for effecting male contraception or other forms of hormone replacement therapy.

5 The AR also regulates female sexual function (i.e., libido), bone formation, and erythropoiesis.

[0003] Worldwide population growth and social awareness of family planning have stimulated a great deal of research in contraception. Contraception is a difficult subject

10 under any circumstances. It is fraught with cultural and social stigma, religious implications, and, most certainly, significant health concerns. This situation is only exacerbated when the subject focuses on male contraception. Despite the availability of suitable contraceptive devices, historically, society has looked to women to be responsible for contraceptive decisions and their consequences. Although health

15 concerns over sexually transmitted diseases have made men more aware of the need to develop safe and responsible sexual habits, women still often bear the brunt of contraceptive choice. Women have a number of choices, from temporary mechanical devices such as sponges and diaphragms to temporary chemical devices such as spermicides. Women also have at their disposal more permanent options, such as

20 physical devices like IUDs and cervical caps as well as more permanent chemical treatments, such as birth control pills and subcutaneous implants. However, to date, the only options available for men include the use of condoms or a vasectomy. Condom use, however is not favored by many men because of the reduced sexual sensitivity, the interruption in sexual spontaneity, and the significant possibility of pregnancy caused by

25 breakage or misuse. Vasectomies are also not favored. If more convenient methods of birth control were available to men, particularly long term methods that require no preparative activity immediately prior to a sexual act, such methods could significantly increase the likelihood that men would take more responsibility for contraception.

30 [0004] Administration of the male sex steroids (e.g., testosterone and its derivatives) has shown particular promise in this regard due to the combined gonadotropin-suppressing and androgen-substituting properties of these compounds (Steinberger et al.,

"Effect of Chronic Administration of Testosterone Enanthate on Sperm Production and Plasma Testosterone, Follicle Stimulating Hormone, and Luteinizing Hormone Levels: A Preliminary Evaluation of a Possible Male Contraceptive", Fertility and Sterility 28:1320- 28 (1977)). Chronic administration of high doses of testosterone completely  
5 abolishes sperm production (azoospermia) or reduces it to a very low level (oligospermia). The degree of spermatogenic suppression necessary to produce infertility is not precisely known. However, a recent report by the World Health Organization showed that weekly intramuscular injections of testosterone enanthate result in  
10 azoospermia or severe oligospermia (i.e., less than 3 million sperm per ml) and infertility in 98% of men receiving therapy (World Health Organization Task Force on Methods Ar Regulation of Male Fertility, "Contraceptive Efficacy of Testosterone-Induced Azoospermia and Oligospermia in Normal Men," Fertility and Sterility 65:821-29 (1996)).

15 [0005] A variety of testosterone esters have been developed that are more slowly absorbed after intramuscular injection and, thus, result in greater androgenic effect. Testosterone enanthate is the most widely used of these esters. While testosterone enanthate has been valuable in terms of establishing the feasibility of hormonal agents for male contraception, it has several drawbacks, including the need for weekly  
20 injections and the presence of supraphysiologic peak levels of testosterone immediately following intramuscular injection (Wu, "Effects of Testosterone Enanthate in Normal Men: Experience From a Multicenter Contraceptive Efficacy Study," Fertility and Sterility 65:626-36 (1996)).


25 [0006] Steroidal ligands which bind the AR and act as androgens (e.g. testosterone enanthate) or as antiandrogens (e.g. cyproterone acetate) have been known for many years and are used clinically (Wu 1988). Although nonsteroidal antiandrogens are in clinical use for hormone-dependent prostate cancer, nonsteroidal androgens have not  
30 been reported. For this reason, research on male contraceptives has focused solely on steroidal compounds.



## SUMMARY OF THE INVENTION

[0007] This invention provides a novel class of tissue-selective androgen receptor targeting agents (ARTA). The agents define a new subclass of compounds which are tissue-selective androgen receptor modulators (SARM), which are useful for oral testosterone replacement therapy, male contraception, maintaining sexual desire in women, osteoporosis, treating prostate cancer and imaging prostate cancer. These agents have an unexpected and tissue-selective in-vivo activity for an androgenic and anabolic activity of a nonsteroidal ligand for the AR. These agents selectively act as partial agonists in some tissues, while acting as full agonists in other tissues, providing a novel and unexpected means for eliciting tissue-selective androgenic or anabolic effects. These agents may be active alone or in combination with progestins or estrogens. The invention further provides a novel class of non-steroidal agonist compounds. The invention further provides compositions containing the selective androgen modulator compounds or the non-steroidal agonist compounds and methods of binding an AR, modulating spermatogenesis, bone formation and/or resorption, treating and imaging prostate cancer, and providing hormonal therapy for androgen-dependent conditions.

[0008] The present invention relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula I:



wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;


Z is NO<sub>2</sub>, CN, COR, COOH or CONHR;

Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;


Q is alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR

wherein R is a alkyl, aryl, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl;


or Q together with the benzene ring to which it is attached is a fused ring system represented by structure A, B or C:



A



B



C

R<sub>1</sub> is CH<sub>3</sub>, CF<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, or CF<sub>2</sub>CF<sub>3</sub>; and


T is OH, OR, -NHCOCH<sub>3</sub>, or NHCOR wherein R is a C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl.

[0009] In one embodiment, Q is in the para position. In another embodiment, X is O.

In another embodiment, Q is in the para position and X is O. In yet another embodiment, Q is para alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR wherein R is a alkyl, aryl, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl.

[00010] The present invention relates to a selective androgen receptor modulator compound having in-vivo androgenic and anabolic activity of a nonsteroidal ligand for

the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula II:



5

wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;

Z is a hydrogen bond acceptor, NO<sub>2</sub>, CN, COR, CONHR;


Y is a lipid soluble group, I, CF<sub>3</sub>, Br, Cl, SnR<sub>3</sub>;

R is an alkyl or aryl group or OH; and

10

Q is acetamido-, trifluoroacetamido-, alkylamines, ether, alkyl, N-sulfonyl, O-sulfonyl, alkylsulfonyl, carbonyl, or a ketone.

[00011] The present invention also relates to a selective androgen receptor modulator compound having in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor the, selective androgen receptor modulator compound represented by the structure of formula III:



where

X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;

Z is NO<sub>2</sub>, CN, COR, or CONHR;

Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;


20

R is an alkyl, or aryl group or OH; and

Q is acetamido or trifluoroacetamido.

[00012] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a


nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula IV:



IV


5

[00013] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula V:




V

[00014] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula VI:



VI

[00015] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula VII:



VII

[00016] The present invention also relates to a method of binding a selective androgen receptor modulator compound to an androgen receptor, which includes contacting the androgen receptor with the selective androgen receptor modulator compound under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00017] Another aspect of the present invention relates to a method of modulating spermatogenesis in a subject, which includes contacting an androgen receptor of the subject with a selective androgen receptor modulator compound under conditions effective to increase or decrease sperm production. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another

embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00018] The present invention also relates to a method of hormone therapy, comprising  
5 contacting an androgen receptor of a subject with a selective androgen receptor  
modulator compound under conditions effective to bind the selective androgen receptor  
modulator compound to the androgen receptor and effect a change in an androgen-  
dependent condition. In one embodiment the compound is Compound I. In another  
embodiment the compound is Compound II. In another embodiment the compound is  
10 Compound III. In another embodiment the compound is Compound IV. In another  
embodiment the compound is Compound V. In another embodiment the compound is  
Compound VI. In another embodiment the compound is Compound VII. In another  
embodiment the compound is Compound VIII.

15 [00019] The present invention also relates to a method of treating a subject having a  
hormone related condition which comprises contacting an androgen receptor of said  
subject with a selective androgen receptor modulator compound under conditions  
effective to bind the selective androgen receptor modulator compound to the androgen  
receptor and effect a change in an androgen-dependent condition. In one embodiment,  
20 the selective androgen receptor modulator compound is selective for androgen or  
testosterone receptor. The present invention also relates to a method of oral  
administration of the selective androgen receptor modulator compound.

[00020] The present invention also relates to a method of treating a subject having a  
25 chronic muscle wasting disease state which comprises contacting an androgen receptor  
of said subject with a selective androgen receptor modulator compound as described  
herein under conditions effective to bind the selective androgen receptor modulator  
compound to the androgen receptor and effect a change in an androgen-dependent  
condition. In one embodiment, the selective androgen receptor modulator compound is  
30 selective for androgen or testosterone receptor. The present invention also relates to a  
method of oral administration of the selective androgen receptor modulator compound.  
In one embodiment the compound is Compound I. In another embodiment the


compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00021] As defined herein, the disease state of "chronic muscle wasting" means

[00022] The present invention also relates to a method of treating a subject having prostate cancer which comprises administering to a subject an effective amount of a selective androgen receptor modulator compound. In one embodiment, the selective androgen receptor modulator compound is selective for androgen or testosterone receptor. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00023] The present invention also relates to compositions and a pharmaceutical compositions which comprises a selective androgen receptor modulator alone or in combination with a progestin or estrogen and a suitable carrier, diluent or salt. In one embodiment the composition comprises Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.


[00024] The present invention relates to a non-steroidal agonist compound, the non-steroidal agonist compound represented by the structure of formula VIII:



wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;  
 R<sub>1</sub> is CH<sub>3</sub>, CF<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, or CF<sub>2</sub>CF<sub>3</sub>;  
 T is OH, OR, -NHCOCH<sub>3</sub>, or NHCOR wherein R is a C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl;


5

A is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or heterocyclic ring represented by the structure:



10

B is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or heterocyclic ring represented by the structure:



15

wherein A<sub>1</sub>-A<sub>11</sub> are each C, O, S or N;  
 B<sub>1</sub>-B<sub>11</sub> are each C, O, S or N;  
 Z is NO<sub>2</sub>, CN, COOH COR, or CONHR;  
 Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>; and

20

Q<sub>1</sub> and Q<sub>2</sub> are independently of each other alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR wherein R is a C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl.

25



- [00025] The present invention also relates to a composition and pharmaceutical composition comprising the non-steroidal agonist compound alone or in combination with a progestin or estrogen and a suitable carrier, diluent or salt. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.
- 10 [00026] The present invention also relates to a method of binding a non-steroidal agonist compound to an androgen receptor comprising contacting the androgen receptor with the non-steroidal agonist compound under conditions effective to bind the non-steroidal agonist compound to the androgen receptor. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.
- 15 [00027] The present invention also relates to a method of modulating spermatogenesis in a subject comprising contacting an androgen receptor of the subject with a non-steroidal agonist compound under conditions effective to increase or decrease sperm production. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.
- 20 [00028] The present invention also relates to a method of hormone therapy comprising contacting an androgen receptor of a subject with a non-steroidal agonist under conditions effective to bind the non-steroidal agonist compound to the androgen receptor
- 25
- 30

and effect a change in an androgen-dependent condition. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In  
5 another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00029] The present invention also relates to a method of treating a subject having a hormone related condition which comprises contacting an androgen receptor of said  
10 subject with a non-steroidal agonist compound under conditions effective to bind the non-steroidal agonist compound to the androgen receptor and effect a change in an androgen-dependent condition. In one embodiment, the non-steroidal agonist compound is selective for androgen or testosterone receptor. The present invention also relates to a method of oral administration of the non-steroidal agonist compound. In one  
15 embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is  
20 Compound VIII.

[00030] The present invention also relates to a method of treating a subject having prostate cancer which comprises administering to a subject an effective amount of a non-steroidal agonist compound. In one embodiment, the non-steroidal agonist  
25 compound is selective for androgen or testosterone receptor. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the  
30 compound is Compound VII. In another embodiment the compound is Compound VIII.

[00031] Still another aspect of the present relates to a method of producing a selective

androgen receptor modulator or a non-steroidal AR agonist compound of the present invention. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00032] The present invention further relates to a method of determining the presence of a selective androgen modulator compound and/or a non-steroidal agonist compound of the present invention in a sample. The method comprises the steps of obtaining the sample, and detecting the compound in the sample, thereby determining the presence of the compound in the sample. In one embodiment, the sample is a blood serum, plasma, urine, or saliva sample. In another embodiment, the detection step comprises measuring the absorbance of the compound. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00033] The novel selective androgen receptor modulator compounds and the non-steroidal agonist compounds of the present invention, either alone or as a composition, are useful in males and females for the treatment of a variety of hormone-related conditions, such as hypogonadism, sarcopenia, erythropoiesis, erectile function, lack of libido, osteoporosis and fertility. Further, the selective androgen receptor modulator compounds and the non-steroidal agonist compounds are useful for oral testosterone replacement therapy, treating prostate cancer, imaging prostate cancer, and maintaining sexual desire in women. The agents may be used alone or in combination with a progestin or estrogen. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another

embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

5 [00034] The selective androgen receptor modulator compounds and the non-steroidal agonist compounds of the present invention offer a significant advance over steroidal androgen treatment because the selective androgen receptor modulator compounds and the non-steroidal agonist compounds of the present invention have been shown in-vivo to have a tissue-selective androgenic and anabolic activity of a nonsteroidal ligand for  
10 the androgen receptor. Moreover, the selective androgen receptor modulator compounds and the non-steroidal agonist compounds of the present invention are not accompanied by serious side effects, lability to oxidative metabolism, inconvenient modes of administration, or high costs and still have the advantages of oral bioavailability, lack of cross-reactivity with other steroid receptors, and long biological half-lives. In one  
15 embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is  
20 Compound VIII.

### BRIEF DESCRIPTION OF THE DRAWINGS

25 The present invention will be understood and appreciated more fully from the following detailed description taken in conjunction with the appended drawings in which:

30 **Figure 1:** Androgenic and Anabolic activity of (S)-GTx-007 in rats. Rats were left untreated (intact control), castrated (castrated control), treated with testosterone propionate (TP), or treated with S-GTx-007, and the body weight gain as well as the weight of androgen-responsive tissues (prostate, seminal vesicles and levator ani muscle) was determined.

5 **Figure 2:** Androgenic and Anabolic activity of S-GTx-007 in rats. Rats were left untreated (intact control), castrated (castrated control), treated with 0.1, 0.3, 0.5, 0.75 and 1.0 mg/day testosterone propionate (TP), or treated with 0.1, 0.3, 0.5, 0.75 and 1.0 mg/day S-GTx-007, and the weight of androgen-responsive tissues (prostate, seminal vesicles and levator ani muscle) was determined.

10 **Figure 3:** Androgenic and Anabolic activity of S-GTx-014 in rats. Rats were left untreated (intact control), castrated (castrated control), treated with 0.1, 0.3, 0.5, 0.75 and 1.0 mg/day testosterone propionate (TP), or treated with 0.1, 0.3, 0.5, 0.75 and 1.0 mg/day S-GTx-014, and the weight of androgen-responsive tissues (prostate, seminal vesicles and levator ani muscle) was determined.

15 **Figure 4:** Average plasma concentration-time profiles of S-GTx-007 in beagle dogs after IV administration at 3 and 10 mg/kg.

**Figure 5:** Average plasma concentration-time profiles of S-GTx-007 in beagle dogs after PO administration as solution at 10 mg/kg.

20 **Fig 6:** Average plasma concentration-time profiles of S-GTx-007 in beagle dogs after IV administration as capsules at mg/kg.

**Figure 7:** Effects of GTx-014 and GTx-007 on LH Levels.

25 **Figure 8:** Effects of GTx-014 and GTx-007 on FSH Levels.

**Figure 9:** Synthesis scheme of GTX-007.

### DETAILED DESCRIPTION OF THE INVENTION


[00035] This invention provides a novel class of androgen receptor targeting agents (ARTA). The agents define a new subclass of compounds which are tissue-selective androgen receptor modulators (SARM) which are useful for oral testosterone replacement therapy, male contraception, maintaining sexual desire in women, treating prostate cancer and imaging prostate cancer. These agents have an unexpected tissue-selective in-vivo activity for an androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor. These agents may be active alone or in combination with progestins or estrogens. The invention further provides a novel class of non-steroidal agonist compounds. The invention further provides compositions containing the selective androgen modulator compounds or the non-steroidal agonist compounds and methods of binding an androgen receptor, modulating spermatogenesis, treating and imaging prostate cancer, and providing hormonal therapy for androgen-dependent conditions.

[00036] The compounds described herein, define a new class of selective androgen receptor modulators (SARMS) that demonstrate potent anabolic effects (e.g., muscle growth) with less androgenic activity (e.g., prostatic growth). This new class of drugs has several advantages over non-selective androgens, including potential therapeutic applications in males and females for modulation of fertility, erythropoiesis, osteoporosis, sexual libido and in men with or at high risk for prostate cancer. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00037] Further, in one embodiment the compounds have tissue specific pharmacologic activity. As demonstrated in Figure 7 and 8, GTX-007 does not suppress LH levels at doses that are capable of eliciting maximal stimulation of levator ani muscle growth and

does not suppress FSH levels at doses that are capable of eliciting maximal stimulation of levator ani muscle growth.

[00038] The present invention relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor selective androgen receptor modulator compound represented by the structure of formula I:




wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;


Z is NO<sub>2</sub>, CN, COR, COOH or CONHR;

Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;


Q is alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR wherein R is an aryl, C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl; or Q together with the benzene ring to which it is attached is a fused ring system represented by structure A, B or C:



A



B




C

$R_1$  is  $\text{CH}_3$ ,  $\text{CF}_3$ ,  $\text{CH}_2\text{CH}_3$ , or  $\text{CF}_2\text{CF}_3$ ; and

T is OH, OR,  $-\text{NHCOCH}_3$ , or  $\text{NHCOR}$  wherein R is a  $\text{C}_1$ - $\text{C}_4$  alkyl, a  $\text{C}_1$ - $\text{C}_4$  haloalkyl, phenyl, halo, alkenyl or hydroxyl.

[00039] In one embodiment, Q is in the para position of the benzene ring to which it is attached. In another embodiment, Q is in the para position and X is O. In another embodiment, Q is in the para position and is alkyl, halogen,  $\text{NR}_2$ ,  $\text{NHCOCH}_3$ ,  $\text{NHCOCF}_3$ ,  $\text{NHCOR}$ ,  $\text{NHCONHR}$ ,  $\text{NHCOOR}$ ,  $\text{OCONHR}$ ,  $\text{CONHR}$ ,  $\text{NHCSCH}_3$ ,  $\text{NHCSCH}_3$ ,  $\text{NHCSR}$ ,  $\text{NHSO}_2\text{CH}_3$ ,  $\text{NHSO}_2\text{R}$ , OR, COR, OCOR,  $\text{OSO}_2\text{R}$ ,  $\text{SO}_2\text{R}$  or SR wherein R is a, aryl,  $\text{C}_1$ - $\text{C}_4$  alkyl, a  $\text{C}_1$ - $\text{C}_4$  haloalkyl, phenyl, halo, alkenyl or hydroxyl.

[00040] The present invention relates to a selective androgen receptor modulator compound having in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula II:



wherein X is a O,  $\text{CH}_2$ , NH, Se, PR, or NR;

Z is a hydrogen bond acceptor,  $\text{NO}_2$ , CN, COR, CONHR;


Y is a lipid soluble group, I,  $\text{CF}_3$ , Br, Cl,  $\text{SnR}_3$ ;

R is an alkyl, or aryl group or OH; and

Q is acetamido-, trifluoroacetamido-, alkylamines, ether, alkyl, N-sulfonyl, O-sulfonyl, alkylsulfonyl, carbonyl, or a ketone.



[00041] The present invention also relates to a selective androgen receptor modulator compound having in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor the, selective androgen receptor modulator compound represented by the structure of formula III:



where

X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;


Z is NO<sub>2</sub>, CN, COR, or CONHR;

Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;


R is an alkyl or aryl group or OH; and

Q is acetamido or trifluoroacetamido.

[00042] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula IV:




[00043] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula V:




V

[00044] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula VI:




VI

[00045] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula VII:




VII


[00046] The present invention relates to a non-steroidal agonist compound having the formula (Compound VIII):



- 5        wherein    X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;  
                      R<sub>1</sub> is CH<sub>3</sub>, CF<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, or CF<sub>2</sub>CF<sub>3</sub>;  
                      T is OH, OR, -NHCOCH<sub>3</sub>, or NHCOR wherein R is a C<sub>1</sub>-C<sub>4</sub>  
                      alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl;  
                      A is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or  
 10        heterocyclic ring represented by the structure:



- B is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or  
 15        heterocyclic ring represented by the structure:



wherein

A<sub>1</sub>-A<sub>11</sub> are each C, O, S or N;

B<sub>1</sub>-B<sub>11</sub> are each C, O, S or N;

Z is NO<sub>2</sub>, CN, COR, COOH, or CONHR;

Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>; and

Q<sub>1</sub> and Q<sub>2</sub> are independently of each other alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR wherein R is a C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl.

[00047] The substituents Z and Y can be in any position of the five or 6 membered ring carrying these substituents (hereinafter "A ring"). Similarly, the substituent Q can be in any position of the five or 6 membered ring carrying this substituent (hereinafter "B ring"). It is understood that when any of the ring members A<sub>1</sub>-A<sub>11</sub> or B<sub>1</sub>-B<sub>11</sub> are O or S, then these ring members are unsubstituted. It is further understood that when any of the ring members A<sub>1</sub>-A<sub>11</sub> or B<sub>1</sub>-B<sub>11</sub> are O or S, then the dotted line between said ring members and other ring members represents a single bond.

[00048] In one embodiment, the A ring includes any type of saturated or unsaturated carbocyclic ring. In one embodiment, the A ring is a 6 membered saturated carbocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In one embodiment, the A ring is a 5 membered saturated carbocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 6 membered carbocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 5 membered carbocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove.

[00049] In another embodiment, the A ring includes any type of saturated, unsaturated or aromatic heterocyclic ring. In another embodiment, the A ring is a 6 membered saturated heterocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 5 membered saturated heterocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 6 membered heterocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 5 membered heterocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 6 membered heteroaromatic ring which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 5 membered heteroaromatic ring which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove.

[00050] Similarly, the B ring includes any type of saturated or unsaturated carbocyclic ring. In one embodiment, the B ring is a 6 membered saturated carbocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In one embodiment, the B ring is a 5 membered saturated carbocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the B ring is a 6 membered carbocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the B ring is a 5 membered carbocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove.

[00051] In another embodiment, the B ring includes any type of saturated, unsaturated or aromatic heterocyclic ring. In another embodiment, the B ring is a 6 membered saturated heterocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the B ring is a 5 membered saturated heterocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the B ring is a 6 membered heterocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the B ring is a 5 membered heterocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the B ring is a 6 membered heteroaromatic ring which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the B ring is a 5 membered heteroaromatic ring which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove.

Nonlimiting examples of suitable A rings and/or B rings are carbocyclic rings such as cyclopentane, cyclopentene, cyclohexane, and cyclohexene rings, and heterocyclic rings such as pyran, dihydropyran, tetrahydropyran, pyrrole, dihydropyrrole, tetrahydropyrrole, pyrazine, dihydropyrazine, tetrahydropyrazine, pyrimidine, dihydropyrimidine, tetrahydropyrimidine, pyrazol, dihydropyrazol, tetrahydropyrazol, piperidine, piperazine, pyridine, dihydropyridine, tetrahydropyridine, morpholine, thiomorpholine, furan, dihydrofuran, tetrahydrofuran, thiophene, dihydrothiophene, tetrahydrothiophene, thiazole, imidazole, isoxazole, and the like.

[00052] As used herein, receptors for extracellular signaling molecules are collectively referred to as "cell signaling receptors". Many cell signaling receptors are transmembrane proteins on a cell surface; when they bind an extracellular signaling molecule (i.e., a ligand), they become activated so as to generate a cascade of intracellular signals that alter the behavior of the cell. In contrast, in some cases, the

receptors are inside the cell and the signaling ligand has to enter the cell to activate them; these signaling molecules therefore must be sufficiently small and hydrophobic to diffuse across the plasma membrane of the cell. As used herein, these receptors are collectively referred to as "intracellular cell signaling receptors".

5

[00053] Steroid hormones are one example of small hydrophobic molecules that diffuse directly across or are transported across the plasma membrane of target cells and bind to intracellular cell signaling receptors. These receptors are structurally related and constitute the intracellular receptor superfamily (or steroid-hormone receptor superfamily). Steroid hormone receptors include progesterone receptors, estrogen receptors, androgen receptors, glucocorticoid receptors, and mineralocorticoid, and numerous orphan receptors. The present invention is particularly directed to androgen receptors and all of its isoforms.

15

[00054] In addition to ligand binding to the receptors, the receptors can be blocked to prevent ligand binding. When a substance binds to a receptor, the three-dimensional structure of the substance fits into a space created by the three-dimensional structure of the receptor in a ball and socket configuration.

20

[00055] The better the ball fits into the socket, the more tightly it is held. This phenomenon is called affinity. If the affinity of a substance is sufficiently high, it will compete with the hormone and bind the binding site more frequently. The binding of the ligand may also lead to tissue-selective recruitment of other important proteins to transduce the signal. These proteins are known as coactivators and corepressor, participate in signal transduction, and may be selectively induced or inhibited by ligand binding. Once bound, signals may be sent through the receptor into the cells, causing the cell to respond in some fashion. This is called activation. On activation, the activated receptor then directly regulates the transcription of specific genes. But the substance and the receptor may have certain attributes, other than affinity, that activate the cell.

25

30 Chemical bonds between atoms of the substance and the atoms of the receptors may form. In some cases, this leads to a change in the configuration of the receptor, which is enough to begin the activation process (called signal transduction). As a result,

substances can be made which bind receptors and activate them (called receptor agonists) or inactivate them (called receptor antagonists).

[00056] The present invention is directed to selective androgen receptor modulator compounds which are agonist compounds, and are, therefore, useful in binding to and activating steroidal hormone receptors. The compounds are non-steroidal. Preferably, the agonist compound of the present invention is an agonist that binds the androgen receptor. Preferably, the compound has high affinity for the androgen receptor. The compound may bind either reversibly or irreversibly to the androgen receptor. The compound of the present invention may contain a functional group (affinity label) that allows alkylation of the androgen receptor (i.e. covalent bond formation). Thus, in this case, the compound binds irreversibly to the receptor and, accordingly, cannot be displaced by a steroid, such as the endogenous ligands dihydrotestosterone and testosterone. It is preferable, however, for the compounds of the present invention to reversibly bind the androgen receptor.

[00057] According to one aspect of the present invention, a method is provided for binding the selective androgen receptor modulator compounds of the present invention to an androgen receptor by contacting the receptor with a selective androgen receptor modulator compound under conditions effective to cause the selective androgen receptor modulator compound to bind the androgen receptor. The binding of the selective androgen receptor modulator compounds to the androgen receptor enables the compounds of the present invention to be useful in males and in females in a number of hormone therapies. The agonist compounds bind to and activate the androgen receptor. Binding of the agonist compound is either reversible or irreversible, preferably reversible.

[00058] According to one aspect of the present invention, a method is provided for modulating spermatogenesis by contacting an androgen receptor of a patient with a selective androgen receptor modulator compound under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor and increase or decrease sperm production.



[00059] According to another aspect of the present invention, a method is provided for hormonal therapy in a patient (i.e., suffering from an androgen- dependent condition) which includes contacting an androgen receptor of a patient with a selective androgen receptor modulator compound under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor and effect a change in an androgen-dependent condition. Androgen-dependent conditions that may be treated according to the present invention include those conditions associated with aging, such as hypogonadism, sarcopenia, erythropoiesis, osteoporosis, and any other conditions later determined to be dependent upon low androgen (e.g., testosterone) levels. In one embodiment, the selective androgen receptor modulator compound is administered alone. In another embodiment, the selective androgen receptor modulator compound is administered in combination with progestin. In yet another embodiment, the selective androgen receptor modulator compound is administered in combination with estrogen.

[00060] According to another aspect of the present invention, a method is provided for treating a subject having prostate cancer. The method comprises administering to a subject an effective amount of a selective androgen receptor modulator compound. In one embodiment, the selective androgen receptor modulator compound is selective for androgen or testosterone receptor.

[00061] The present invention also relates to a method of treating a subject having a chronic muscle wasting disease state which comprises contacting an androgen receptor of said subject with a selective androgen receptor modulator compound under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor and effect a change in an androgen-dependent condition. In one embodiment, the selective androgen receptor modulator compound is selective for androgen or testosterone receptor. The present invention also relates to a method of oral administration of the selective androgen receptor modulator compound. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is

Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

5

[00062] According to one aspect of the present invention, a method is provided for binding the non-steroidal agonist compounds of the present invention to an androgen receptor by contacting the receptor with a non-steroidal agonist compound under conditions effective to cause the non-steroidal agonist compound to bind the androgen  
10 receptor. The binding of the non-steroidal agonist compounds to the androgen receptor enables the compounds of the present invention to be useful in males and in females in a number of hormone therapies. The agonist compounds bind to and activate the androgen receptor. Binding of the agonist compound is either reversible or irreversible, preferably reversible.

15

[00063] According to one aspect of the present invention, a method is provided for modulating spermatogenesis by contacting an androgen receptor of a patient with a non-steroidal agonist compound under conditions effective to bind the selective androgen  
20 receptor modulator compound to the androgen receptor and increase or decrease sperm production.

[00064] According to another aspect of the present invention, a method is provided for hormonal therapy in a patient (i.e., one suffering from an androgen- dependent condition) which includes contacting an androgen receptor of a patient with a non-steroidal agonist compound under conditions effective to bind the non-steroidal agonist  
25 compound to the androgen receptor and effect a change in an androgen-dependent condition. Androgen-dependent conditions that may be treated according to the present invention include those conditions associated with aging, such as hypogonadism, sarcopenia, erythropoiesis, osteoporosis, lack of sexual libido and any other conditions  
30 later determined to be dependent upon low androgen (e.g., testosterone) levels. In one embodiment, the non-steroidal agonist compound is administered alone. In another embodiment, the non-steroidal agonist compound is administered in combination with

progestin. In yet another embodiment, the non-steroidal agonist compound is administered in combination with estrogen.

[00065] According to another aspect of the present invention, a method is provided for  
5 treating a subject having prostate cancer. The method comprises administering to a  
subject an effective amount of a non-steroidal agonist compound. In one embodiment,  
the non-steroidal agonist compound is selective for androgen or testosterone receptor.

[00066] The compounds of the present invention have an asymmetric center and can be  
10 the R or S isomer, or a mixture of both. In one embodiment, the compounds are racemic  
mixtures of the R and S enantiomers. In another embodiment, the compounds are  
substantially pure R enantiomers. In another embodiment, the compounds are  
substantially pure S enantiomers. "Substantially pure" is defined herein as greater than  
about 95% preponderance of one isomer. Where the above-described processes for the  
15 preparation of the compounds of use in the invention give rise to mixtures of  
stereoisomers, these isomers may be separated by conventional techniques, such as  
preparative chromatography. The compounds may be prepared in racemic form, or  
individual enantiomers may be prepared either by enantiospecific synthesis or by  
resolution

20  
[00067] As used herein, "pharmaceutical composition" means therapeutically effective  
amounts of the SARM or the non-steroidal agonist compound of the present invention,  
together with suitable diluents, preservatives, solubilizers, emulsifiers, adjuvant and/or  
carriers. A "therapeutically effective amount" as used herein refers to that amount which  
25 provides a therapeutic effect for a given condition and administration regimen. Such  
compositions are liquids or Lyophilized or otherwise dried formulations and include  
diluents of various buffer content (e.g., Tris-HCl, acetate, phosphate), pH and ionic  
strength, additives such as albumin or gelatin to prevent absorption to surfaces,  
detergents (e.g., Tween 20, Tween 80, Pluronic F68, bile acid salts), solubilizing agents  
30 (e.g., glycerol, polyethylene glycerol), anti-oxidants (e.g., ascorbic acid, sodium  
metabisulfite), preservatives (e.g., Thimerosal, benzyl alcohol, parabens), bulking  
substances or tonicity modifiers (e.g., lactose, mannitol), covalent attachment of

polymers such as polyethylene glycol to the protein, complexation with metal ions, or incorporation of the material into or onto particulate preparations of polymeric compounds such as polylactic acid, polglycolic acid, hydrogels, etc, or onto liposomes, microemulsions, micelles, unilamellar or multilamellar vesicles, erythrocyte ghosts, or spheroplasts. Such compositions will influence the physical state, solubility, stability, rate of in vivo release, and rate of in vivo clearance. Controlled or sustained release compositions include formulation in lipophilic depots (e.g., fatty acids, waxes, oils).

[00068] Also comprehended by the invention are particulate compositions coated with polymers (e.g., poloxamers or poloxamines). Other embodiments of the compositions of the invention incorporate particulate forms, protective coatings, protease inhibitors or permeation enhancers for various routes of administration, including parenteral, pulmonary, nasal and oral. In one embodiment the pharmaceutical composition is administered parenterally, paracancerally, transmucosally, transdermally, intramuscularly, intravenously, intradermally, subcutaneously, intraperitoneally, intraventricularly, intracranially and intratumorally.

[00069] Further, as used herein "pharmaceutically acceptable carriers" are well known to those skilled in the art and include, but are not limited to, 0.01-0.1M and preferably 0.05M phosphate buffer or 0.8% saline. Additionally, such pharmaceutically acceptable carriers may be aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media.

[00070] Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers such as those based on Ringer's dextrose, and the like. Preservatives and other additives may also be present, such as, for example, antimicrobials, antioxidants, collating agents, inert gases and the like.

[00071] Controlled or sustained release compositions include formulation in lipophilic depots (e.g. fatty acids, waxes, oils). Also comprehended by the invention are particulate compositions coated with polymers (e.g. poloxamers or poloxamines) and the compound  
5 coupled to antibodies directed against tissue-specific receptors, ligands or antigens or coupled to ligands of tissue-specific receptors.

[00072] Other embodiments of the compositions of the invention incorporate particulate forms, protective coatings, protease inhibitors or permeation enhancers for  
10 various routes of administration, including parenteral, pulmonary, nasal and oral.

[00073] Compounds modified by the covalent attachment of water-soluble polymers such as polyethylene glycol, copolymers of polyethylene glycol and polypropylene glycol, carboxymethyl cellulose, dextran, polyvinyl alcohol, polyvinylpyrrolidone or  
15 polyproline are known to exhibit substantially longer half-lives in blood following intravenous injection than do the corresponding unmodified compounds (Abuchowski et al., 1981; Newmark et al., 1982; and Katre et al., 1987). Such modifications may also increase the compound's solubility in aqueous solution, eliminate aggregation, enhance the physical and chemical stability of the compound, and greatly reduce the  
20 immunogenicity and reactivity of the compound. As a result, the desired in vivo biological activity may be achieved by the administration of such polymer-compound adducts less frequently or in lower doses than with the unmodified compound.

[00074] In yet another embodiment, the pharmaceutical composition can be delivered  
25 in a controlled release system. For example, the agent may be administered using intravenous infusion, an implantable osmotic pump, a transdermal patch, liposomes, or other modes of administration. In one embodiment, a pump may be used (see Langer, supra; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989). In another embodiment,  
30 polymeric materials can be used. In yet another embodiment, a controlled release system can be placed in proximity to the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled

Release, supra, vol. 2, pp. 115-138 (1984). Preferably, a controlled release device is introduced into a subject in proximity to the site of inappropriate immune activation or a tumor. Other controlled release systems are discussed in the review by Langer (Science 249:1527-1533 (1990)).

5

[00075] The pharmaceutical preparation can comprise the selective androgen receptor modulator alone, or can further include a pharmaceutically acceptable carrier, and can be in solid or liquid form such as tablets, powders, capsules, pellets, solutions, suspensions, elixirs, emulsions, gels, creams, or suppositories, including rectal and urethral  
10 suppositories. Pharmaceutically acceptable carriers include gums, starches, sugars, cellulosic materials, and mixtures thereof. The pharmaceutical preparation containing the selective androgen receptor modulator can be administered to a subject by, for example, subcutaneous implantation of a pellet; in a further embodiment, the pellet provides for controlled release of selective androgen receptor modulator over a period of  
15 time. The preparation can also be administered by intravenous, intraarterial, or intramuscular injection of a liquid preparation, oral administration of a liquid or solid preparation, or by topical application. Administration can also be accomplished by use of a rectal suppository or a urethral suppository.

20 [00076] The pharmaceutical preparations of the invention can be prepared by known dissolving, mixing, granulating, or tablet-forming processes. For oral administration, the selective androgen receptor modulators or their physiologically tolerated derivatives such as salts, esters, N-oxides, and the like are mixed with additives customary for this purpose, such as vehicles, stabilizers, or inert diluents, and converted by customary  
25 methods into a suitable form for administration, such as tablets, coated tablets, hard or soft gelatin capsules, aqueous, alcoholic or oily solutions. Examples of suitable inert vehicles are conventional tablet bases such as lactose, sucrose, or cornstarch in combination with binders like acacia, cornstarch, gelatin, or with disintegrating agents such as cornstarch, potato starch, alginic acid, or with a lubricant like stearic acid or  
30 magnesium stearate.

[00077] Examples of suitable oily vehicles or solvents are vegetable or animal oils

such as sunflower oil or fish-liver oil. Preparations can be effected both as dry and as wet granules. For parenteral administration (subcutaneous, intravenous, intraarterial, or intramuscular injection), the SARM agents or the non-steroidal agonist agents or their physiologically tolerated derivatives such as salts, esters, N-oxides, and the like are converted into a solution, suspension, or emulsion, if desired with the substances customary and suitable for this purpose, for example, solubilizers or other auxiliaries. Examples are: sterile liquids such as water and oils, with or without the addition of a surfactant and other pharmaceutically acceptable adjuvants. Illustrative oils are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, or mineral oil. In general, water, saline, aqueous dextrose and related sugar solutions, and glycols such as propylene glycols or polyethylene glycol are preferred liquid carriers, particularly for injectable solutions.

[00078] The preparation of pharmaceutical compositions which contain an active component is well understood in the art. Typically, such compositions are prepared as aerosols of the polypeptide delivered to the nasopharynx or as injectables, either as liquid solutions or suspensions, however, solid forms suitable for solution in, or suspension in, liquid prior to injection can also be prepared. The preparation can also be emulsified. The active therapeutic ingredient is often mixed with excipients that are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof.

[00079] In addition, if desired, the composition can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, which enhance the effectiveness of the active ingredient.

[00080] An active component can be formulated into the composition as neutralized pharmaceutically acceptable salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide or antibody molecule), which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like.

Salts formed from the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

5

[00081] For topical administration to body surfaces using, for example, creams, gels, drops, and the like, the SARM agents or the non-agonist steroidal compounds or their physiologically tolerated derivatives such as salts, esters, N-oxides, and the like are prepared and applied as solutions, suspensions, or emulsions in a physiologically acceptable diluent with or without a pharmaceutical carrier.

10

[00082] In another embodiment, the active compound can be delivered in a vesicle, in particular a liposome (see Langer, Science 249:1527-1533 (1990); Treat et al., in Liposomes in the Therapy of Infectious Disease and Cancer, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*).

15

[00083] For use in medicine, the salts of the SARM or the non-steroidal agonist compounds will be pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds according to the invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts, which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, methanesulphonic acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, oxalic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid.

20

25

[00084] The present invention further relates to a method of determining the presence of a selective androgen modulator compound and/or a non-steroidal agonist compound of the present invention in a sample. The method comprises the steps of obtaining the sample, and detecting the compound in the sample, thereby determining the presence of the compound in the sample.

30



[00085] In one embodiment, the sample is a blood serum sample. In another embodiment, the sample is a plasma sample. In another embodiment, the sample is a urine sample. In another embodiment, the sample is a saliva sample. In another  
5 embodiment, the sample is any other tissue sample.

[00086] In one embodiment, the detection step comprises measuring the absorbance of the compound at a predetermined wavelength. For example, the compounds of the present invention absorb in the ultraviolet region of the spectrum, with an absorbency  
10 peak at 270 nm. Thus, in one embodiment of the present invention, the compound is detected by monitoring the UV absorbance of the sample at 270 nm. It should be noted that the present invention is not limited to UV absorption, and that any other spectrometric methods of identification are applicable. For example, compounds can be detected by measuring their infra-red or visible absorbance.

15 [00087] In another embodiment, the present invention further provides a method of determining the concentration of a selective androgen receptor modulator compound and/or a non-steroidal agonist compound of the present invention in a sample. The method comprises the steps of obtaining a sample; determining the level of the  
20 compound in the sample, and calculating the concentration of the compound in the sample by comparing the level with a standard sample containing a known concentration of the compound. Calibration curves of known concentrations of the compound in the sample, can be obtained, and the concentration of the compound in the test sample is calculated therefrom. By "level" it is meant the absorption level of the compound at the  
25 measured wavelength.

[00088] In another embodiment, the compound is detected in the sample by contacting the sample with a binding protein which specifically binds to the compound, and determining the amount of binding protein bound to the compound. The concentration  
30 of the compound can be determined by measuring the amount of binding protein bound to the compound, and comparing that amount to a standard sample containing a known concentration of the compound - binding protein complex.

5 [00089] Protein levels can be determined according to standard techniques, as described in Sambrook et al. Briefly, a sample obtained from a subject is contacted with a binding protein which specifically binds to a specific compound of the present invention, and the amount of complex formed between the binding protein and the compound is determined. In one embodiment, the binding protein is an antibody which specifically binds to one or more compounds of the present invention. In another embodiment, the binding protein has a detectable label bound thereto, and the complex between the binding protein-label compound is determined by visualizing the complex.

10 [00090] As defined herein, "contacting" means that the binding protein is introduced into the sample in a test tube, flask, tissue culture, chip, array, plate, microplate, capillary, or the like, and incubated at a temperature and time sufficient to permit the binding component to bind to a cell or a fraction thereof or plasma/serum or a fraction thereof containing the target. Methods for contacting the samples with the binding proteins, or other specific binding components are known to those skilled in the art and may be selected depending on the type of assay protocol to be run. Incubation methods are also standard and are known to those skilled in the art.

20 [00091] "Visualizing" the complex may be carried out by any means known in the art, including, but not limited to, ELISA, radioimmunoassay, flow cytometry, dot blots, western immunoblotting combined with gel electrophoresis, immunohistochemistry at light and electron pe levels, HPLC and mass spectrometry.

25 [00092] Either monoclonal or polyclonal antibodies (as well as any recombinant antibodies) specific for the selective androgen modulator compounds or the non-steroidal agonist compounds of the present invention can be used in the various immunoassays. The antibodies may be detectably labeled, utilizing conventional labeling techniques well-known to the art. As used herein, the term "label" refers to a molecule, which may be conjugated or otherwise attached (i.e., covalently or non-covalently) to a binding protein as defined herein. Labels are known to those skilled in the art. Thus, the antibodies may be labeled with radioactive isotopes, non-radioactive

30

isotopic labels, fluorescent labels, enzyme labels, chemiluminescent labels, bioluminescent labels, free radical labels, or bacteriophage labels, using techniques known in the art. Examples of radioisotopic labels are <sup>3</sup>H, <sup>125</sup>I, <sup>131</sup>I, <sup>35</sup>S, <sup>14</sup>C, etc. Examples of non-radioactive isotopic labels are <sup>55</sup>Mn, <sup>56</sup>Fe, etc. Examples of fluorescence labels are fluorescent labels which are directly labeled with the preferred fluorescence label, or fluorescent labels which are indirectly labeled with the preferred fluorescence label. In the last case, the preferred fluorescence label is conjugated to a secondary antibody, which is directed against the first antibody, such as an anti species Ig antibody. Typical fluorescent labels include, but are not limited to a fluorescein label, an isothiocyanate label, a rhodamine label, a phycoerythrin label, etc., for example fluorescein isothiocyanate (FITC, International Biological Supplies, Melbourne, FL), rhodamine, phycoerythrin (P.E., Coulter Corp., Hialeah, FL), phycocyanin, allophycocyanin, phycoerythrin-cyanin dye 5 (PECy5, Coulter), label, a phycocyanin label, an allophycocyanin label, an O-phthaldehyde label, a fluorescamine and Texas Red.

[00093] Examples of enzyme labels include alkaline phosphatase, beta-galactosidase, glucose-6-phosphate dehydrogenase, maleate dehydrogenase, and peroxidase. Two principal types of enzyme immunoassay are the enzyme-linked immunosorbent assay (ELISA), and the homogeneous enzyme immunoassay, also known as enzyme-multiplied immunoassay (EMIT, Syva Corporation, Palo Alto, CA). In the ELISA system, separation may be achieved, for example, by the use of antibodies coupled to a solid phase. The EMIT system depends on deactivation of the enzyme in the tracer-antibody complex; the activity can thus be measured without the need for a separation step.

[00094] Particularly suitable labels include those, which permit analysis by flow cytometry, e.g., fluorochromes. Other suitable detectable labels include those useful in colorimetric enzyme systems, e. g., horseradish peroxidase (HRP) and alkaline phosphatase (AP). Other proximal enzyme systems are known to those of skill in the art, including hexokinase in conjunction with glucose-6-phosphate dehydrogenase.

[00095] Additionally, chemiluminescent compounds may be used as labels. Chemiluminescent labels, such as green fluorescent proteins, blue fluorescent proteins, and variants thereof are known. Also bioluminescence or chemiluminescence can be detected using, respectively, NAD oxidoreductase with luciferase and substrates NADH and FNIN or peroxidase with luminol and substrate peroxide. Typical chemiluminescent compounds include luminol, isoluminol, aromatic acridinium esters, imidazoles, acridinium salts, and oxalate esters. Similarly, bioluminescent compounds may be utilized for labelling, the bioluminescent compounds including luciferin, luciferase, and aequorin. Once labeled, the antibody may be employed to identify and quantify immunologic counterparts (antibody or antigenic polypeptide) utilizing techniques well-known to the art.

[00096] The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way be construed, however, as limiting the broad scope of the invention.

## EXPERIMENTAL DETAILS SECTION

### EXAMPLE 1

#### **Nonsteroidal Ligands with Androgenic and Anabolic Activity**

[00097] The SARM compounds provided herein were designed, synthesized and evaluated for in-vitro and in-vivo pharmacologic activity. The in-vitro androgen receptor binding affinity and ability to maintain androgen dependent tissue growth in castrated animals was studied. Androgenic activity was monitored as the ability of the SARM compounds to maintain and/or stimulate the growth of the prostate and seminal vesicles, as measured by weight. Anabolic activity was monitored as the ability of the SARM compounds to maintain and/or stimulate the growth of the levator ani muscle, as measured by weight.

Synthetic Procedures of Compounds I-VIII:

[00098] (2R)-1-Methacryloylpyrrolidin-2-carboxylic Acid (R-129). D-Proline (R-128, 14.93 g, 0.13 mol) was dissolved in 71 mL of 2 N NaOH and cooled in an ice bath; the resulting alkaline solution was diluted with acetone (71 mL). An acetone solution (71 mL) of metacryloyl chloride 127 (13.56 g, 0.13 mol) and 2N NaOH solution (71 mL) were simultaneously added over 40 min to the aqueous solution of D-proline in an ice bath. The pH of the mixture was kept at 10-11°C during the addition of the metacryloyl chloride. After stirring (3 h, room temperature), the mixture was evaporated in vacuo at a temperature at 35-45 °C to remove acetone. The resulting solution was washed with ethyl ether and was acidified to pH 2 with concentrated HCl. The acidic mixture was saturated with NaCl and was extracted with EtOAc (100 mL x 3). The combined extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered through Celite, and evaporated in vacuo to give the crude product as a colorless oil. Recrystallization of the oil from ethyl ether and hexanes afforded 16.2 (68%) of the desired compound as colorless crystals: mp 102-103 °C (lit. [214] mp 102.5-103.5 °C); the NMR spectrum of this compound demonstrated the existence of two rotamers of the title compound. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 5.28 (s) and 5.15 (s) for the first rotamer, 5.15 (s) and 5.03 (s) for the second rotamer (totally 2H for both rotamers, vinyl CH<sub>2</sub>), 4.48-4.44 for the first rotamer, 4.24-4.20 (m) for the second rotamer (totally 1H for both rotamers, CH at the chiral center), 3.57-3.38 (m, 2H, CH<sub>2</sub>), 2.27-2.12 (1H, CH), 1.97-1.72 (m, 6H, CH<sub>2</sub>, CH, Me); <sup>13</sup>C NMR (75 MHz, DMSO-d<sub>6</sub>) δ for major rotamer 173.3, 169.1, 140.9, 116.4, 58.3, 48.7, 28.9, 24.7, 19.5; for minor rotamer 174.0, 170.0, 141.6, 115.2, 60.3, 45.9, 31.0, 22.3, 19.7; IR (KBr) 3437 (OH), 1737 (C=O), 1647 (CO, COOH), 1584, 1508, 1459, 1369, 1348, 1178 cm<sup>-1</sup>; [α]<sub>D</sub><sup>26</sup> +80.8° (c = 1, MeOH); Anal. Calcd. for C<sub>9</sub>H<sub>13</sub>NO<sub>3</sub>: C 59.00, H 7.15, N 7.65. Found: C 59.13, H 7.19, N 7.61.

[00099] (3R,8aR)-3-Bromomethyl-3-methyl-tetrahydro-pyrrolo[2,1-c][1,4]oxazine-1,4-dione (R, R-130). A solution of NBS (23.5g, 0.132 mol) in 100 mL of DMF was added dropwise to a stirred solution of compound R-129 (16.1g, 88 mmol) in 70 mL of DMF under argon at room temperature, and the resulting mixture was stirred 3 days. The solvent was removed in vacuo, and a yellow solid was precipitated.

The solid was suspended in water, stirred overnight at room temperature, filtered, and dried to give 18.6 (81%) (smaller weight when dried ~ 34%) of the title compound as a yellow solid: mp 152-154 °C (lit. [214] mp 107-109 °C for the S-isomer); <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 4.69 (dd, J = 9.6 Hz, J = 6.7 Hz, 1H, CH at the chiral center), 4.02 (d, J = 11.4 Hz, 1H, CHH<sub>a</sub>), 3.86 (d, J = 11.4 Hz, 1H, CHH<sub>b</sub>), 3.53-3.24 (m, 4H, CH<sub>2</sub>), 2.30-2.20 (m, 1H, CH), 2.04-1.72 (m, 3H, CH<sub>2</sub> and CH), 1.56 (s, 2H, Me); <sup>13</sup>C NMR (75 MHz, DMSO-d<sub>6</sub>) δ 167.3, 163.1, 83.9, 57.2, 45.4, 37.8, 29.0, 22.9, 21.6; IR (KBr) 3474, 1745 (C=O), 1687 (C=O), 1448, 1377, 1360, 1308, 1227, 1159, 1062 cm<sup>-1</sup>; [α]<sub>D</sub><sup>26</sup> +124.5° (c = 1.3, chloroform); Anal. Calcd. for C<sub>9</sub>H<sub>12</sub>BrNO<sub>3</sub>: C 41.24, H 4.61, N 5.34. Found: C 41.46, H 4.64, N 5.32.

[000100] (2R)-3-Bromo-2-hydroxy-2-methylpropanoic Acid (R-131). A mixture of bromolactone R-130 (18.5g, 71 mmol) in 300 mL of 24% HBr was heated at reflux for 1 h. The resulting solution was diluted with brine (200 mL), and was extracted with ethyl acetate (100 mL x 4). The combined extracts were washed with saturated NaHCO<sub>3</sub> (100 mL x 4). The aqueous solution was acidified with concentrated HCl to pH = 1, which, in turn, was extracted with ethyl acetate (100 mL x 4). The combined organic solution was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered through Celite, and evaporated in vacuo to dryness. Recrystallization from toluene afforded 10.2 g (86%) of the desired compound as colorless crystals: mp 107-109 °C (lit. [214] mp 109-113 °C for the S-isomer); <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 3.63 (d, J = 10.1 Hz, 1H, CHH<sub>a</sub>), 3.52 (d, J = 10.1 Hz, 1H, CHH<sub>b</sub>), 1.35 (s, 3H, Me); IR (KBr) 3434 (OH), 3300-2500 (COOH), 1730 (C=O), 1449, 1421, 1380, 1292, 1193, 1085 cm<sup>-1</sup>; [α]<sub>D</sub><sup>26</sup> +10.5° (c = 2.6, MeOH); Anal. Calcd. for C<sub>4</sub>H<sub>7</sub>BrO<sub>3</sub>: C 26.25, H 3.86. Found: C 26.28, H 3.75.

[000101] N-[4-Nitro-3-(trifluoromethyl)phenyl]-(2R)-3-bromo-2-hydroxy-2-methylpropanamide (R-132). Thionyl chloride (8.6 g, 72 mmol) was added dropwise under argon to a solution of bromoacid R-131 (11.0 g, 60 mmol) in 70 mL of DMA at -5 to -10 °C. The resulting mixture was stirred for 2 h under the same conditions. A solution of 4-nitro-3-trifluoromethyl-aniline (12.4 g, 60 mmol) in 80 mL of DMA was added dropwise to the above solution, and the resulting mixture was stirred overnight at room temperature. The solvent was removed on Rotavapor using high vacuum oil pump;

the residue was diluted with saturated NaHCO<sub>3</sub> solution, and extracted with ethyl ether (100 mL x 3). Combined extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered through Celite, and purified by flash chromatography on silica gel, using methylene chloride as eluent to afford 18.0 g (80%) of the desired compound: mp 98-100 °C (*R*<sub>f</sub> = 0.2, silica gel, CH<sub>2</sub>Cl<sub>2</sub>); <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 10.54 (s, 1H, NH), 8.54 (d, *J* = 2.1 Hz, 1H, ArH), 8.34 (dd, *J* = 9.0 Hz, *J* = 2.1 Hz, 1H, ArH), 8.18 (d, *J* = 9.0 Hz, 1H, ArH), 6.37 (s, 1H, OH), 3.82 (d, *J* = 10.4 Hz, 1H, CHH<sub>a</sub>), 3.58 (d, *J* = 10.4 Hz, 1H, CHH<sub>b</sub>), 1.48 (s, 3H, Me); <sup>13</sup>C NMR (75 MHz, DMSO-d<sub>6</sub>) δ 173.6 (C=O), 143.0, 127.2, 123.2, 122.6 (q, *J* = 33.0 Hz), 122.0 (q, *J* = 271.5 Hz), 118.3 (q, *J* = 6.0 Hz), 74.4, 41.4, 24.9; IR (KBr) 3344 (OH), 1680 (C=O), 1599, 1548 (C=C, Ar), 1427, 1363, 1161 cm<sup>-1</sup>; MS (ESI): *m/z* 370.8 (M)<sup>+</sup>; Anal. Calcd. for C<sub>11</sub>H<sub>10</sub>BrN<sub>2</sub>O<sub>4</sub>: C 35.60, H 2.72, N 7.55. Found: C 35.68, H 2.72, N 7.49.

[000102] N-[4-nitro-3-trifluoromethyl)phenyl]-(2S)-3-[4-(acetylamino)phenoxy]-2-hydroxy-2-methylpropanamide (S-147). The title compound was prepared from compound R-132 (0.37 g, 1.0 mmol), 4-acetamidophenol (0.23 g, 1.5 mmol) K<sub>2</sub>CO<sub>3</sub> (0.28 g, 2.0 mmol), and 10% of benzyltributylammonium chloride as a phase transfer catalyst in 20 mL of methyl ethyl ketone was heated at reflux overnight under argon. The reaction was followed by TLC, the resulting mixture was filtered through Celite, and concentrated in vacuo to dryness. Purification by flash column chromatography on silica gel (hexanes-ethyl acetate, 3:1) yielded 0.38 g (86%) (*R*<sub>f</sub> = 0.18 hexanes-ethyl acetate, 3:1) of the desired compound as a light yellow powder: mp 70-74 °C; The solid can be recrystallized from ethyl acetate and hexane); <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 10.62 (s, 1H, NH), 9.75 (s, 1H, NH), 8.56 (d, *J* = 1.9 Hz, 1H, ArH), 8.36 (dd, *J* = 9.1 Hz, *J* = 1.9 Hz, 1H, ArH), 8.18 (d, *J* = 9.1 Hz, 1H, ArH), 7.45-7.42 (m, 2H, ArH), 6.85-6.82 (m, 2H, ArH), 6.25 (s, 1H, OH), 4.17 (d, *J* = 9.5 Hz, 1H, CHH<sub>a</sub>), 3.94 (d, *J* = 9.5 Hz, 1H, CHH<sub>b</sub>), 1.98 (s, 3H, Me), 1.43 (s, 3H, Me); <sup>13</sup>C NMR (75 MHz, DMSO-d<sub>6</sub>) δ 174.6 (C=O), 167.7, 154.2, 143.3, 141.6, 132.8, 127.4, 123.0, 122.7 (q, *J* = 33.0 Hz), 122.1 (q, *J* = 271.5 Hz), 120.1, 118.3 (q, *J* = 6.0 Hz), 114.6, 74.9, 73.8, 23.8, 23.0; IR (KBr) 3364 (OH), 1668 (C=O), 1599, 1512 (C=C, Ar), 1457, 1415, 1351, 1323, 1239, 1150 1046 cm<sup>-1</sup>; MS (ESI): *m/z* 464.1 (M+Na)<sup>+</sup>; Anal. Calcd. for C<sub>19</sub>H<sub>18</sub>F<sub>3</sub>N<sub>3</sub>O<sub>6</sub>: C 51.71, H 4.11, N 9.52. Found: C 52.33, H 4.40, N 9.01.

[Duane insert]

5

[000103] The in-vitro activity of the SARM compounds, specifically compound VII,  
10 demonstrated high androgen receptor binding affinity ( $K_i = 7.5$  nM). Animal studies  
with the SARM compounds, specifically compound V, demonstrated that it is a potent  
androgenic and anabolic nonsteroidal agent. Four groups of rats were used for these  
studies: (1) intact controls, (2) castrated controls, (3) castrated animals treated with  
testosterone propionate (100  $\mu$ g/day), and (4) castrated animals treated with compound  
15 V (1000  $\mu$ g/day). Testosterone and compound VII were delivered at a constant rate for  
14 days via subcutaneous osmotic pumps.

[000104] The results of these studies are shown in Figure 1. Castration significantly  
reduced the weight of androgenic (e.g., prostate and seminal vesicles) and anabolic (e.g.,  
20 levator ani muscle) tissues, but had little effect on animal body weight (BW). Treatment  
of castrated animals with testosterone propionate or compound VII maintained the  
weight of androgenic tissues to the same degree. Compound VII had similar androgenic  
activity as testosterone propionate (i.e., the prostate and seminal vesicle weights were  
the same), but much greater efficacy as an anabolic agent. Compound VII showed  
25 greater anabolic activity than testosterone propionate at the doses tested (i.e., the levator  
ani muscle maintained the same weight as intact control animals and was greater than  
that observed for testosterone). The experiments presented herein are the first in-vivo  
results which demonstrate tissue-selective androgenic and anabolic activity (i.e.,  
differing androgenic and anabolic potency) of a nonsteroidal ligand for the androgen  
30 receptor.




**EXAMPLE 2****Nonsteroidal Ligands with Androgenic and Anabolic Activity**

5 [000105] The in-vivo efficacy and acute toxicity of four novel nonsteroidal androgens (compounds IV, V, VI and VII) in rats was examined. In-vitro assays established that these compounds bind the androgen receptor with very high affinity. The structures and names of the four compounds are presented below:

10

15



GTx-014	R= F
GTx-015	R=COCH <sub>3</sub>
GTx-016	R=COC <sub>2</sub> H <sub>5</sub>
GTx-017	R=NHCOCH <sub>3</sub>

20

**EXPERIMENTAL METHODS**

25 [000106] *Materials.* The S-isomers of compounds GTx-014 (compound IV), GTx-015 (compound V), GTx-016 (compound VI) and GTx-007 (compound VII wherein R is NHCOCH<sub>3</sub>) and the R-isomer of GTx-014 were synthesized in accordance with the scheme as set forth in Figure 9. Testosterone propionate (TP), polyethylene glycol 300 (PEG300, reagent grade) and neutral buffered formalin (10% w/v) were purchased from

Sigma Chemical Company (St Louis, MO). Alzet osmotic pumps (model 2002) were purchased from Alza Corp. (Palo Alto, CA).

5 [000107] *Animals.* Immature male Sprague-Dawley rats, weighing 90 to 100g, were purchased from Harlan Biosciences (Indianapolis, IN). The animals were maintained on a 12-hour light-dark cycle with food and water available *ad libitum*. The animal protocol was reviewed and approved by the Institutional Laboratory Animal Care and Use Committee.

10 [000108] *Study Design.* Rats were randomly distributed into twenty-nine (29) groups, with 5 animals per group. Treatment groups are described in Table 1. One day prior to the start of drug treatment, animals in groups 2 through 29 were individually removed from the cage, weighed and anesthetized with an intraperitoneal dose of ketamine/xylazine (87/13 mg/kg; approximately 1 mL per kg). When appropriately anesthetized (i.e., no  
15 response to toe pinch), the animals' ears were marked for identification purposes. Animals were then placed on a sterile pad and their abdomen and scrotum washed with betadine and 70% alcohol. The testes were removed via a midline scrotal incision, with sterile suture being used to ligate supra-testicular tissue prior to surgical removal of each testis. The surgical wound site was closed with sterile stainless steel wound clips, and  
20 the site cleaned with betadine. The animals were allowed to recover on a sterile pad (until able to stand) and then returned to their cage.

[000109] Twenty-four hours later, animals in groups 2 through 29 were re-anesthetized with ketamine/xylazine, and an Alzet osmotic pump(s) (model 2002) was placed  
25 subcutaneously in the scapular region. In this instance, the scapular region was shaved and cleaned (betadine and alcohol) and a small incision (1 cm) made using a sterile scalpel. The osmotic pump was inserted and the wound closed with a sterile stainless steel wound clip. Animals were allowed to recover and were returned to their cage. Osmotic pumps contained the appropriate treatment (designated in Table 1) dissolved in  
30 polyethylene glycol 300 (PEG300). Osmotic pumps were filled with the appropriate solution one day prior to implantation. Animals were monitored daily for signs of acute toxicity to drug treatment (e.g., lethargy, rough coat).

[000110] After 14 days of drug treatment, rats were anesthetized with ketamine/xylazine. Animals were then sacrificed by exsanguinations under anesthesia. A blood sample was collected by venipuncture of the abdominal aorta, and submitted for complete blood cell analysis. A portion of the blood was placed in a separate tube,  
5 centrifuged at 12,000g for 1 minute, and the plasma layer removed and frozen at -20°C. The ventral prostates, seminal vesicles, levator ani muscle, liver, kidneys, spleen, lungs, and heart were removed, cleared of extraneous tissue, weighed, and placed in vials containing 10% neutral buffered formalin. Preserved tissues were sent to GTx, Inc. for histopathological analysis.

10

[000111] For data analysis, the weights of all organs were normalized to body weight, and analyzed for any statistical significant difference by single-factor ANOVA. The weights of prostate and seminal vesicle were used as indexes for evaluation of androgenic activity, and the levator ani muscle weight was used to evaluate the anabolic  
15 activity.

### RESULTS

[000112] The androgenic and anabolic activities the S isomers of compounds GTx-014, GTx-015, GTx-016 and GTx-007, and the R isomer of GTx-014 were examined in a  
20 castrated rat model after 14 days of administration. Testosterone propionate, at increasing doses, was used as the positive control of anabolic and androgenic effects.

[000113] As shown in Figures 2 and 3, the weights of prostate, seminal vesicle, and levator ani muscle in castrated, vehicle-treated rats decreased significantly, due to the  
25 ablation of endogenous androgen production. Exogenous administration of testosterone propionate, an androgenic and anabolic steroid, increased the weights of prostate, seminal vesicle, and levator ani muscle in castrated rats in a dose-dependent manner. The R-isomer of GTx-014, and S-isomers of GTx-015 and GTx-016 showed no effect on the weights of prostate, seminal vesicle, and levator ani muscle in castrated animals  
30 (data not shown). The S-isomers of GTx-007 (Figure 2: S-GTx-007) and GTx-014 (Figure 3: S-GTx-014) resulted in dose-dependent increases in prostate, seminal vesicle and levator ani muscle weights. Compared with testosterone propionate, S-GTx-007

showed lower potency and intrinsic activity in increasing the weights of prostate and seminal vesicle, but a greater potency and intrinsic activity in increasing the weight of levator ani muscle. Particularly, S-GTx-007, at a dose as low as 0.3 mg/day, was able to maintain the levator ani muscle weight of castrated animals in the same level as that of intact animals. Thus, S-GTx-007 is a potent nonsteroidal anabolic agent with less androgenic activity but more anabolic activity than testosterone propionate. This is a significant improvement over previous claims, in that this compound selectively stimulates muscle growth and other anabolic effects while having less effect on the prostate and seminal vesicles. This may be particularly relevant in aging men with concerns related to the development or progression of prostate cancer.

[000114] GTx-014 was less potent than GTx-007, but showed greater tissue selectivity (compare effects on the prostate and seminal vesicles in Figures 2 and 3). GTx-014 significantly increased levator ani muscle weights, but showed little to no ability to stimulate prostate and seminal vesicle growth (i.e., the prostate and seminal vesicle weights were less than 20% of that observed in intact animals or in animals treated with testosterone propionate).

[000115] Results showed that none of the examined compounds produced significant effect on body weight or the weights of other organs (i.e., liver, kidneys, spleen, lungs and heart). Nor did any compound produce any signs of acute toxicity, as gauged by diagnostic hematology tests and visual examination of animals receiving treatments. Importantly, GTx-007 did not suppress the production of luteinizing hormone (LH) or follicle stimulating hormone (FSH) at a dose of 0.3 mg/day (i.e., a dose that exhibited maximal anabolic effects).

[000116] In summary, S-GTx-007 exhibited exceptional anabolic activity in animals by maintaining the weight of levator ani muscle after removal of endogenous androgen. This discovery represents major progress towards the development of therapeutically useful nonsteroidal androgens, and a major improvement (i.e., tissue selectivity and potency) over previous drugs in this class. S-GTx-014 and S-GTx-007 showed selective anabolic activity in comparison with testosterone propionate, an androgenic and anabolic

steroid. The tissue-selective activity is actually one of the advantages of nonsteroidal androgens in terms of anabolic-related applications.

5 [000117] Despite similarities in structure and in-vitro functional activity, the S-isomers of compounds GTx-014, GTx-015, GTx-016, and GTx-007 exhibited profound differences in terms of their in-vivo activity. GTx-007 the most efficacious androgenic and anabolic activity in animals, with the anabolic activity greater than that of testosterone propionate. GTx-014 showed a small degree of androgenic activity, but an anabolic activity comparable to testosterone propionate. In contrast, GTx-015 and GTx-016 failed to produce any androgenic or anabolic activity in-vivo.

15 [000118] These studies show the discovery of two members (GTx-014 and GTx-007, compounds, compounds II and V respectively) of a new class of selective androgen receptor modulators (SARMS) that demonstrate potent anabolic effects (e.g., muscle growth) with less androgenic activity (e.g., prostatic growth). This new class of drugs has several advantages over non-selective androgens, including potential therapeutic applications in males and females for modulation of fertility, erythropoiesis, osteoporosis, sexual libido and in men with or at high risk for prostate cancer.

20 [000119] Further, Figures 7 and 8 demonstrate the effects of GTx-014 and GTx-007 on LH and FSH levels in rats. These results further demonstrate the novelty of these SARMS, due to their differential effects on these reproductive hormones, thus demonstrating the tissue-specific pharmacologic activity. In Figure 7, LH levels in castrated animals treated with TP and GTx-014 were significantly lower than those of untreated animals (i.e., castrated controls) at doses greater than or equal to 0.3 mg/day. However, higher doses (i.e., 0.5 mg/day or higher) of GTx-007 were required before significant decreases in LH levels were observed. Thus, GTx-007 does not suppress LH levels at doses that are capable of eliciting maximal stimulation of levator ani muscle growth. In Figure 8, FSH levels in castrated animals treated with GTx-014 were significantly lower than those of untreated animals (i.e., castrated controls) at doses of 0.5 mg/day or higher. Similarly, lower FSH levels were observed in animals treated with TP. However, only this difference was only significant at a dose of 0.75 mg/day. FSH

levels in animals treated with GTx-007 were not significantly different from those of untreated animals at any dose level tested. Thus, GTx-007 does not suppress FSH levels at doses that are capable of eliciting maximal stimulation of levator ani muscle growth.

5 **Table 1. Animals Groups and Experimental Design**

Group #	Castrated?	Drug	Dose	# of animals
1	No	None	None	5
2	Yes	None	Vehicle only	5
3	Yes	Testosterone	0.1 mg/day	5
4	Yes	Testosterone	0.3 mg/day	5
5	Yes	Testosterone	0.5 mg/day	5
6	Yes	Testosterone	0.75 mg/day	5
7	Yes	Testosterone	1.0 mg/day	5
8	Yes	R-GTx-014	1.0 mg/day	5
9	Yes	S-GTx-014	0.1 mg/day	5
10	Yes	S-GTx-014	0.3 mg/day	5
11	Yes	S-GTx-014	0.5 mg/day	5
12	Yes	S-GTx-014	0.75 mg/day	5
13	Yes	S-GTx-014	1.0 mg/day	5
14	Yes	S-GTx-015	0.1 mg/day	5
15	Yes	S-GTx-015	0.3 mg/day	5
16	Yes	S-GTx-015	0.5 mg/day	5
17	Yes	S-GTx-015	0.75 mg/day	5
18	Yes	S-GTx-015	1.0 mg/day	5
19	Yes	S-GTx-016	0.1 mg/day	5
20	Yes	S-GTx-016	0.3 mg/day	5
21	Yes	S-GTx-016	0.5 mg/day	5
22	Yes	S-GTx-016	0.75 mg/day	5
23	Yes	S-GTx-016	1.0 mg/day	5
24	Yes	S-GTx-007	0.1 mg/day	5
25	Yes	S-GTx-007	0.3 mg/day	5
26	Yes	S-GTx-007	0.5 mg/day	5
27	Yes	S-GTx-007	0.75 mg/day	5
28	Yes	S-GTx-007	1.0 mg/day	5
29	Yes	None	Vehicle only	5

5

**EXAMPLE 3****Pharmacokinetics of GTx-007 in Dogs**

10 [000120] The pharmacokinetics of S-GTx-007, a novel selective androgen receptor modulator (SARM), were characterized in beagle dogs. A four-treatment, four-period crossover design was utilized in the study, which involved a total of six beagle dogs, three of each gender. Each animal received a 3 mg/kg IV dose, a 10 mg/kg IV dose, a 10 mg/kg PO dose in solution, and a 10 mg/kg PO dose in capsule, in a randomly assigned  
15 order. There was an one-week washout period between treatments. Plasma samples were collected for up to 72 hr after drug administration. Plasma S-GTx-007 concentrations were analyzed by a validated HPLC method. The clearance (CL), volume of distribution (Vss), half-life ( $T_{1/2}$ ), and other pharmacokinetic parameters were determined by noncompartmental methods. Results showed that S-GTx-007 was cleared from dog  
20 plasma with a terminal  $T_{1/2}$  of about 4 hr and a CL of 4.4 mL/min/kg after IV administration. Figures 4, 5, and 6 show the plasma concentration-time profiles of S-GTx-007 after administration of an intravenous solution, oral solution, and oral capsule, respectively. The pharmacokinetics were dose- and gender-independent. The oral bioavailability of S-GTx-007 varied with the dosage form, and averaged 38% and 19%  
25 for solution and capsule, respectively. Thus, S-GTx-007 demonstrated moderate half-life, slow clearance and moderate bioavailability in beagle dogs, identifying it as the first of a new class of orally bioavailable tissue-selective androgen receptor modulators.

**EXAMPLE 4****GTx-007 Analysis by HPLC**

30 [000121] A reversed phase high pressure liquid chromatograph (HPLC) assay was

developed to quantitate GTx-007 concentrations in dog plasma. Dog blood samples were obtained by venipuncture and centrifuged at 1000g for 15 minutes. Samples were stored frozen at -20°C until analysis. Individual samples were rapidly thawed and an aliquot (0.5 ml) was spiked with internal standard (20 µl of a 200 µg/ml aqueous solution of CM-II-87). An aliquot of 1 ml of acetonitrile was added to the samples to precipitate plasma proteins. The samples were vortexed and then centrifuged at 1000g for 15 minutes. The supernatant was decanted into glass extraction tubes and 7.5 ml of ethyl acetate was added. The extraction mixture was left at room temperature for 20 minutes, and vortexed several times during this interval. The samples were then centrifuged at 1000g for 10 minutes, and the organic phase was removed and placed in conical-bottomed glass tubes. The organic phase was evaporated under nitrogen. The samples were reconstituted in 200 µl of mobile phase (35:65 acetonitrile:water) and transferred to an autosampler vial for HPLC injection (Waters 717 plus autosampler, Waters Corp., Milford, MA). The isocratic mobile phase of 35% (v/v) acetonitrile in water was pumped at a flow rate of 1 ml/min (Model 510, Waters Corp.). The stationary phase was a C18 reversed phase column (Novapak C18, 3.9 x 150 mm). Analytes were monitored with UV detection at 270 nm (Model 486 absorbance detector, Waters Corp.). Retention times for GTx-007 and CM-II-87 were 11.1 and 16.9 minutes, respectively. Chromatography data was collected and analyzed using Millennium software. Plasma concentrations of GTx-007 in each sample were determined by comparison to calibration curves. Calibration curves were constructed by adding known amounts of GTx-007 to dog plasma. Final GTx-007 concentrations in dog plasma samples used in the calibration curves were 0.08, 0.2, 0.4, 2, 4, 10, and 20 µg/ml. Calibration curves were linear over this concentration range and exhibited correlation coefficients ( $r^2$ ) of 0.9935 or greater. Intra- and inter-day coefficients of variation for the standards ranged from 6.4% for 0.08 µg/ml to 7.9% for 20 µg/ml.


[000122] Melting points were determined on a Thomas-Hoover capillary melting point apparatus and are uncorrected. Infrared spectra were recorded on a Perkin Elmer System 2000 FT-IR. Optical rotations were determined on an Autopol<sup>®</sup> III Automatic Polarimeter (Rudolph Research Model III-589-10, Fairfield, New Jersey). Proton and carbon-13 magnetic resonance spectra were obtained on a Bruker AX 300 spectrometer



(300 and 75 MHz for  $^1\text{H}$  and  $^{13}\text{C}$ , respectively). Chemical shift values were reported as parts per million ( $\delta$ ) relative to tetramethylsilane (TMS). Spectral data were consistent with assigned structures. Mass spectra were determined on a Bruker-HP Esquire LC System. Elemental analyses were performed by Atlantic Microlab Inc. (Norcross, GA), and found values were within 0.4 % of the theoretical values. Routine thin-layer chromatography (TLC) was performed on silica gel on aluminum plates (silica gel 60 F 254, 20 x 20 cm, Aldrich Chemical Company Inc., Milwaukee, WI). Flash chromatography was performed on silica gel (Merck, grade 60, 230-400 mesh, 60). Tetrahydrofuran (THF) was dried by distillation over sodium metal. Acetonitrile (MeCN) and methylene chloride ( $\text{CH}_2\text{Cl}_2$ ) were dried by distillation from  $\text{P}_2\text{O}_5$ .

WHAT IS CLAIMED IS:

1. A selective androgen receptor modulator compound having tissue-selective in-  
 vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor  
 5 represented by the structure of formula I:




wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;


Z is NO<sub>2</sub>, CN, COR, COOH or CONHR;

10 Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;


Q is alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHCSO<sub>2</sub>CH<sub>3</sub>, NHCSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR wherein R is an alkyl, aryl, C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl; or Q together  
 15 with the benzene ring to which it is attached is a fused ring system represented by structure A, B or C:



A



B



C

$R_1$  is  $\text{CH}_3$ ,  $\text{CF}_3$ ,  $\text{CH}_2\text{CH}_3$ , or  $\text{CF}_2\text{CF}_3$ ; and

T is OH, OR,  $-\text{NHCOCH}_3$ , or  $\text{NHCOR}$  wherein R is a  $\text{C}_1$ - $\text{C}_4$  alkyl, a  $\text{C}_1$ - $\text{C}_4$  haloalkyl, phenyl, halo, alkenyl or hydroxyl.

2. The selective androgen receptor modulator compound of claim 1, wherein Q is in the para position.


3. The selective androgen receptor modulator compound, of claim 2, wherein Q is alkyl, halogen,  $\text{NR}_2$ ,  $\text{NHCOCH}_3$ ,  $\text{NHCOCF}_3$ ,  $\text{NHCOR}$ ,  $\text{NHCONHR}$ ,  $\text{NHCOOR}$ ,  $\text{OCONHR}$ ,  $\text{CONHR}$ ,  $\text{NHCSCH}_3$ ,  $\text{NHCSCF}_3$ ,  $\text{NHCSR N HSO}_2\text{CH}_3$ ,  $\text{NHSO}_2\text{R}$ , OR, COR, OCOR,  $\text{OSO}_2\text{R}$ ,  $\text{SO}_2\text{R}$  or SR wherein R is an alkyl, aryl,  $\text{C}_1$ - $\text{C}_4$  alkyl, a  $\text{C}_1$ - $\text{C}_4$  haloalkyl, phenyl, halo, alkenyl or hydroxyl.

4. The selective androgen receptor modulator compound of claim 1, wherein X is O.

5. The selective androgen receptor modulator compound of claim 4, wherein Q is in the para position.


6. The selective androgen receptor modulator compound, of claim 5, wherein Q is alkyl, halogen,  $\text{NR}_2$ ,  $\text{NHCOCH}_3$ ,  $\text{NHCOCF}_3$ ,  $\text{NHCOR}$ ,  $\text{NHCONHR}$ ,  $\text{NHCOOR}$ ,  $\text{OCONHR}$ ,  $\text{CONHR}$ ,  $\text{NHCSCH}_3$ ,  $\text{NHCSCF}_3$ ,  $\text{NHCSR N HSO}_2\text{CH}_3$ ,  $\text{NHSO}_2\text{R}$ , OR, COR, OCOR,  $\text{OSO}_2\text{R}$ ,  $\text{SO}_2\text{R}$  or SR wherein R is a  $\text{C}_1$ - $\text{C}_4$  alkyl, a  $\text{C}_1$ - $\text{C}_4$  haloalkyl, phenyl, halo, alkenyl or hydroxyl.

7. The selective androgen receptor modulator compound of claim 1, wherein Z is para NO<sub>2</sub>.
8. The selective androgen receptor modulator compound of claim 1, wherein Y is meta CF<sub>3</sub>.
9. The selective androgen receptor modulator compound of claim 1, wherein T is OH.
10. The selective androgen receptor modulator compound of claim 1, wherein R<sub>1</sub> is CH<sub>3</sub>.
11. The selective androgen receptor modulator compound of claim 1, wherein Q is para F.
12. The selective androgen receptor modulator compound of claim 1, represented by the structure of formula IV.



IV


13. The selective androgen receptor modulator compound of claim 1, wherein Q is para COCH<sub>3</sub>.
14. The selective androgen receptor modulator compound of claim 1, represented by the structure of formula V.



V 55

15. The selective androgen receptor modulator compound of claim 1, wherein Q is para  $\text{COCH}_2\text{CH}_3$ .


5 16. The selective androgen receptor modulator compound of claim 1, represented by the structure of formula VI.



VI

17. The selective androgen receptor modulator compound of claim 1, wherein Q is para  $\text{NHCOCH}_3$ .

10 18. The selective androgen receptor modulator compound of claim 1, represented by the structure of formula VII.



VII

15 19. A composition comprising the selective androgen receptor modulator compound of claim 1 and a suitable carrier or diluent alone or in combination with a progestin or estrogen.

20 20. A pharmaceutical composition comprising an effective amount of the selective androgen receptor modulator compound of claim 1 alone or in combination with a progestin or estrogen and a pharmaceutically acceptable carrier, diluent or salt.

21. A method of binding a selective androgen receptor modulator compound to a androgen receptor comprising contacting the androgen receptor with the selective androgen receptor modulator compound of claim 1 under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor.

5

22. A method of modulating spermatogenesis in a subject comprising contacting an androgen receptor of the subject with the selective androgen receptor modulator compound of claim 1 under conditions effective to increase or decrease sperm production.

10

23. A method of hormone therapy comprising contacting an androgen receptor of a subject with the selective androgen receptor modulator compound of claim 1 under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor and effect a change in an androgen-dependent condition.

15


24. A method of treating a subject having a hormone related condition comprising contacting an androgen receptor of a subject with the selective androgen receptor modulator compound of claim 1 under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor and effect a change in an androgen-dependent condition.

20

25. A method of treating a subject having prostate cancer comprising administering to said subject an effective amount of a selective androgen receptor modulator compound according to claim 1.

25

26. A selective androgen receptor modulator compound having in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor having the formula III:




where X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;  
Z is NO<sub>2</sub>, CN, COR, or CONHR;  
Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;  
R is an alkyl group or OH; and  
Q is acetamido or trifluoroacetamido.

27. The selective androgen receptor modulator compound of claim 26, wherein X is O.
28. The selective androgen receptor modulator compound of claim 26, wherein Z is NO<sub>2</sub>.
29. The selective androgen receptor modulator compound of claim 26, wherein Y is CF<sub>3</sub>.
30. The selective androgen receptor modulator compound of claim 26, wherein Q is NHCOCH<sub>3</sub>.
31. The selective androgen receptor modulator compound of claim 26, wherein X is O, Z is NO<sub>2</sub>, Y is CF<sub>3</sub>, and Q is NHCOCH<sub>3</sub>.
32. A composition comprising the selective androgen receptor modulator compound of claim 26 and a suitable carrier or diluent.
33. A pharmaceutical composition comprising an effective amount of the selective androgen receptor modulator compound of claim 26 and a pharmaceutically acceptable carrier, diluent or salt.
34. A method of binding a selective androgen receptor modulator compound to a androgen receptor comprising contacting the androgen receptor with the selective androgen receptor modulator compound of claim 26 effective to bind the selective androgen receptor modulator compound to the androgen receptor.
35. A method of suppressing spermatogenesis in a subject comprising contacting an androgen receptor of the subject with the selective androgen receptor modulator compound of claim 26 effective to suppress sperm production.
36. A method of hormone therapy comprising contacting an androgen receptor of a subject with the selective androgen receptor modulator compound of claim 26 effective

to bind the selective androgen receptor modulator compound to the androgen receptor and effect a change in an androgen-dependent condition.

37. A method of treating a subject having a hormone related condition comprising contacting an androgen receptor of a subject with the selective androgen receptor modulator compound of claim 26 effective to bind the selective androgen receptor modulator compound to the androgen receptor and effect a change in an androgen-dependent condition.

38. A nonsteroidal agonist compound represented by the structure of formula VIII:



10


wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;

R<sub>1</sub> is CH<sub>3</sub>, CF<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, or CF<sub>2</sub>CF<sub>3</sub>;

T is OH, OR, -NHCOCH<sub>3</sub>, or NHCOR wherein R is a C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl;

15


A is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or heterocyclic ring represented by the structure:



B is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or heterocyclic ring represented by the structure:

20





wherein

A<sub>1</sub>-A<sub>11</sub> are each C, O, S or N;

B<sub>1</sub>-B<sub>11</sub> are each C, O, S or N;

Z is NO<sub>2</sub>, CN, COR, COOH or CONHR;

Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>; and

Q<sub>1</sub> and Q<sub>2</sub> are independently of each other alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR wherein R is a C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl.

39. A composition comprising the non-steroidal agonist compound of claim 38 and a suitable carrier or diluent alone or in combination with a progestin or estrogen.

40. A pharmaceutical composition comprising an effective amount of the non-steroidal agonist compound of claim 38 alone or in combination with a progestin or estrogen and a pharmaceutically acceptable carrier, diluent or salt.

41. A method of binding a compound to an androgen receptor comprising contacting the androgen receptor with the non-steroidal agonist compound of claim 38 under conditions effective to bind the non-steroidal agonist compound to the androgen receptor.

42. A method of modulating spermatogenesis in a subject comprising contacting an androgen receptor of the subject with the non-steroidal agonist compound of claim 38 under conditions effective to increase or decrease sperm production.

43. A method of hormone therapy comprising contacting an androgen receptor of a subject with the non-steroidal agonist compound of claim 38 under conditions effective

to bind the non-steroidal agonist compound to the androgen receptor and effect a change in an androgen-dependent condition.

5 44. A method of treating a subject having a hormone related condition comprising contacting an androgen receptor of a subject with the non-steroidal agonist compound of claim 38 under conditions effective to bind the non-steroidal agonist compound to the androgen receptor and effect a change in an androgen-dependent condition.

10 45. A method of treating a subject having prostate cancer comprising administering to said subject an effective amount of the non-steroidal agonist compound according to claim 38.

15 46. A method of determining the presence of the compound of claim 1 in a sample, said method comprising the steps of:  
obtaining said sample; and  
detecting said compound in said sample;  
thereby determining the presence of said compound in said sample.

20 47. The method according to claim 46, wherein the sample is a blood serum, plasma, urine, or saliva sample.

48. The method according to claim 46, wherein the detection step comprises measuring the absorbance of the compound.

25 49. A method of determining the presence of the compound of claim 26 in a sample, said method comprising the steps of:  
obtaining said sample; and  
detecting said compound in said sample;  
thereby determining the presence of said compound in said sample.

30 50. The method according to claim 49, wherein the sample is a blood serum, plasma, urine, or saliva sample.

51. The method according to claim 49, wherein the detection step comprises measuring the absorbance of the compound.
- 5 52. A method of determining the presence of the compound of claim 38 in a sample, said method comprising the steps of:
- obtaining said sample; and
- detecting said compound in said sample;
- thereby determining the presence of said compound in said sample.
- 10 53. The method according to claim 52, wherein the sample is a blood serum, plasma, urine, or saliva sample.
- 15 54. The method according to claim 52, wherein the detection step comprises measuring the absorbance of the compound.




FIGURE 1

FIGURE 2. Androgenic and anabolic activity of compound II (S-GTx-007) in rats




FIGURE 3. Androgenic and anabolic activity of compound IV (S-GTx-014) in rats





Fig. 4. Average plasma concentration-time profiles of S-GTx-007 in beagle dogs after IV administration at 3 and 10 mg/kg. Each data point represents the mean  $\pm$  standard deviation of 3 animals.



**FIGURE 5.** Average plasma concentration-time profiles of S-GTx-007 in beagle dogs after PO administration as solution at 10 mg/kg. Each data point represents the mean  $\pm$  standard deviation of 3 animals.







FIGURE 6. Plasma concentration-time profiles of S-GTx-007 in beagle dogs after PO administration as capsules at 10 mg/kg.






FIGURE 7







FIGURE 8

# **GTx-007** **Synthetic Outline**



**GTx-007**  
(Acetamidoxolutamide)

Figure 9

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/26328

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : Please See Extra Sheet.

US CL : 514/493, 616, 619; 564/153, 156, 157, 158

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/493, 616, 619; 564/153, 156, 157, 158

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
None

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAPLUS on STN

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4,636,505 A (TUCKER) 13 January 1987 (13/01/87), see entire document.	1-54
A	EP 0 253 503 A2 (IMPERIAL CHEMICAL INDUSTRIES PLC) 20 January 1988 (20/01/88), see entire document.	1-54

☐

Further documents are listed in the continuation of Box C.

☐

See patent family annex.

\*

Special categories of cited documents:

"A"

document defining the general state of the art which is not considered to be of particular relevance

"E"

earlier document published on or after the international filing date

"L"

document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O"

document referring to an oral disclosure, use, exhibition or other means

"P"

document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"A"

document member of the same patent family

Date of the actual completion of the international search

20 SEPTEMBER 2001

Date of mailing of the international search report

16 NOV 2001

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BRIAN J. DAVIS

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/26328

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

C07C 233/15, 233/33, 233/48, 235/26, 237/14; A01N 37/18; A01K 31/16, 31/165

CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 February 2002 (28.02.2002)

PCT

(10) International Publication Number  
**WO 02/016310 A1**

(51) International Patent Classification<sup>7</sup>: **C07C 233/15**,  
233/33, 233/48, 235/26, 237/14, A01N 37/18, 31/16,  
31/165

(21) International Application Number: PCT/US01/26328

(22) International Filing Date: 23 August 2001 (23.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/644,970 24 August 2000 (24.08.2000) US  
60/300,083 25 June 2001 (25.06.2001) US

(71) Applicant (*for all designated States except US*): **GTX, INC.** [US/US]; 3 North Dunlap, Van Fleet Building, Memphis, TN 38163 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **DALTON, James** [US/US]; 2706 Wellesley Drive, Columbus, OH 43221 (US). **MILLER, Duane, D.** [US/US]; 8706 Maple Creek Cove, Germantown, TN 38139 (US). **YIN, Donghua** [US/US]; 984 Chatham Lane, Apt B, Columbus, OH 43221 (US). **HE, Yali** [US/US]; 812-H South Parker Drive, Florence, SC 25901 (US).

(74) Agent: **COHEN, Mark, S.**; Eitan, Pearl, Latzer & Cohen-Zedek, One Crystal Park, Suite 210, 2011 Crystal Drive, Arlington, VA 22202-3709 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(48) Date of publication of this corrected version:

18 July 2002

(15) Information about Correction:

see PCT Gazette No. 29/2002 of 18 July 2002, Section II

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: SELECTIVE ANDROGEN RECEPTOR MODULATORS AND METHODS OF USE THEREOF

(57) Abstract: This invention provides a novel class of androgen receptor targeting agents (ARTA). The agents define a new subclass of compounds which are tissue-selective androgen receptor modulators (SARM), which are useful for oral testosterone replacement therapy, male contraception, maintaining sexual desire in women, treating prostate cancer and imaging prostate cancer. These agents have an unexpected in-vivo activity for an androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor. These agents may be active alone or in combination with progestins or estrogens. The invention further provides a novel class of non-steroidal agonist compounds. The invention further provides compositions containing the selective androgen modulator compounds or the non-steroidal agonist compounds and methods of binding an androgen receptor, modulating spermatogenesis, treating and imaging prostate cancer, and providing hormonal therapy for androgen-dependent conditions.

WO 02/016310 A1

**SELECTIVE ANDROGEN RECEPTOR MODULATORS AND  
METHODS OF USE THEREOF**

**FIELD OF INVENTION**

5

[0001] The present invention relates to a novel class of tissue-selective androgen receptor targeting agents (ARTA) which demonstrate androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor. The agents define a new subclass of compounds which are tissue-selective androgen receptor modulators (SARM) which are  
10 useful for male hormone therapy such as oral testosterone replacement therapy, male contraception, maintaining sexual desire in women, treating prostate cancer, and imaging prostate cancer. These agents are also administered to a subject for the treatment of sarcopenia, lack of sexual libido, osteoporosis, erythropoiesis, and fertility. The agents may be used alone or in combination with a progestin or estrogen.

15

**BACKGROUND OF THE INVENTION**

[0002] The androgen receptor ("AR") is a ligand-activated transcriptional regulatory protein that mediates induction of male sexual development and function through its activity with endogenous androgens. Androgens are generally known as the male sex  
20 hormones. However, androgens also play a pivotal role in female physiology and reproduction. The androgenic hormones are steroids which are produced in the body by the testis and the cortex of the adrenal gland, or synthesized in the laboratory. Androgenic steroids play an important role in many physiologic processes, including the development and maintenance of male sexual characteristics such as muscle and bone  
25 mass, prostate growth, spermatogenesis, and the male hair pattern (Matsumoto, Endocrinol. Met. Clin. N. Am. 23:857-75 (1994)). The endogenous steroidal androgens include testosterone and dihydrotestosterone ("DHT"). Testosterone is the principal steroid secreted by the testes and is the primary circulating androgen found in the plasma of males. Testosterone is converted to DHT by the enzyme 5 alpha-reductase in many  
30 peripheral tissues. DHT is thus thought to serve as the intracellular mediator for most androgen actions (Zhou, et al., Molec. Endocrinol. 9:208-18 (1995)). Other steroidal androgens include esters of testosterone, such as the cypionate, propionate, phenylpropionate, cyclopentylpropionate, isocaproate, enanthate, and decanoate esters, and other synthetic androgens such as 7-Methyl-Nortestosterone ("MENT") and its



acetate ester (Sundaram et al., "7 Alpha-Methyl-Nortestosterone(MENT): The Optimal Androgen For Male Contraception," Ann. Med., 25:199-205 (1993) ("Sundaram")). Because the AR is involved in male sexual development and function, the AR is a likely target for effecting male contraception or other forms of hormone replacement therapy.

5 The AR also regulates female sexual function (i.e., libido), bone formation, and erythropoiesis.

[0003] Worldwide population growth and social awareness of family planning have stimulated a great deal of research in contraception. Contraception is a difficult subject

10 under any circumstances. It is fraught with cultural and social stigma, religious implications, and, most certainly, significant health concerns. This situation is only exacerbated when the subject focuses on male contraception. Despite the availability of suitable contraceptive devices, historically, society has looked to women to be responsible for contraceptive decisions and their consequences. Although health

15 concerns over sexually transmitted diseases have made men more aware of the need to develop safe and responsible sexual habits, women still often bear the brunt of contraceptive choice. Women have a number of choices, from temporary mechanical devices such as sponges and diaphragms to temporary chemical devices such as spermicides. Women also have at their disposal more permanent options, such as

20 physical devices like IUDs and cervical caps as well as more permanent chemical treatments, such as birth control pills and subcutaneous implants. However, to date, the only options available for men include the use of condoms or a vasectomy. Condom use, however is not favored by many men because of the reduced sexual sensitivity, the interruption in sexual spontaneity, and the significant possibility of pregnancy caused by

25 breakage or misuse. Vasectomies are also not favored. If more convenient methods of birth control were available to men, particularly long term methods that require no preparative activity immediately prior to a sexual act, such methods could significantly increase the likelihood that men would take more responsibility for contraception.

30 [0004] Administration of the male sex steroids (e.g., testosterone and its derivatives) has shown particular promise in this regard due to the combined gonadotropin-suppressing and androgen-substituting properties of these compounds (Steinberger et al.,

"Effect of Chronic Administration of Testosterone Enanthate on Sperm Production and Plasma Testosterone, Follicle Stimulating Hormone, and Luteinizing Hormone Levels: A Preliminary Evaluation of a Possible Male Contraceptive", Fertility and Sterility 28:1320- 28 (1977)). Chronic administration of high doses of testosterone completely  
5 abolishes sperm production (azoospermia) or reduces it to a very low level (oligospermia). The degree of spermatogenic suppression necessary to produce infertility is not precisely known. However, a recent report by the World Health Organization showed that weekly intramuscular injections of testosterone enanthate result in azoospermia or severe oligospermia (i.e., less than 3 million sperm per ml) and infertility  
10 in 98% of men receiving therapy (World Health Organization Task Force on Methods Ar Regulation of Male Fertility, "Contraceptive Efficacy of Testosterone-Induced Azoospermia and Oligospermia in Normal Men," Fertility and Sterility 65:821-29 (1996)).


15 [0005] A variety of testosterone esters have been developed that are more slowly absorbed after intramuscular injection and, thus, result in greater androgenic effect. Testosterone enanthate is the most widely used of these esters. While testosterone enanthate has been valuable in terms of establishing the feasibility of hormonal agents for male contraception, it has several drawbacks, including the need for weekly  
20 injections and the presence of supraphysiologic peak levels of testosterone immediately following intramuscular injection (Wu, "Effects of Testosterone Enanthate in Normal Men: Experience From a Multicenter Contraceptive Efficacy Study," Fertility and Sterility 65:626-36 (1996)).

25 [0006] Steroidal ligands which bind the AR and act as androgens (e.g. testosterone enanthate) or as antiandrogens (e.g. cyproterone acetate) have been known for many years and are used clinically (Wu 1988). Although nonsteroidal antiandrogens are in clinical use for hormone-dependent prostate cancer, nonsteroidal androgens have not been reported. For this reason, research on male contraceptives has focused solely on  
30 steroidal compounds.

## SUMMARY OF THE INVENTION

[0007] This invention provides a novel class of tissue-selective androgen receptor targeting agents (ARTA). The agents define a new subclass of compounds which are tissue-selective androgen receptor modulators (SARM), which are useful for oral testosterone replacement therapy, male contraception, maintaining sexual desire in women, osteoporosis, treating prostate cancer and imaging prostate cancer. These agents have an unexpected and tissue-selective in-vivo activity for an androgenic and anabolic activity of a nonsteroidal ligand for the AR. These agents selectively act as partial agonists in some tissues, while acting as full agonists in other tissues, providing a novel and unexpected means for eliciting tissue-selective androgenic or anabolic effects. These agents may be active alone or in combination with progestins or estrogens. The invention further provides a novel class of non-steroidal agonist compounds. The invention further provides compositions containing the selective androgen modulator compounds or the non-steroidal agonist compounds and methods of binding an AR, modulating spermatogenesis, bone formation and/or resorption, treating and imaging prostate cancer, and providing hormonal therapy for androgen-dependent conditions.

[0008] The present invention relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula I:



wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;


Z is NO<sub>2</sub>, CN, COR, COOH or CONHR;

Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;


5 Q is alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR

wherein R is a alkyl, aryl, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl;


10 or Q together with the benzene ring to which it is attached is a fused ring system represented by structure A, B or C:



A



B



C

15 R<sub>1</sub> is CH<sub>3</sub>, CF<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, or CF<sub>2</sub>CF<sub>3</sub>; and


T is OH, OR, -NHCOCH<sub>3</sub>, or NHCOR wherein R is a C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl.

[0009] In one embodiment, Q is in the para position. In another embodiment, X is O.  
 20 In another embodiment, Q is in the para position and X is O. In yet another embodiment, Q is para alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR wherein R is a alkyl, aryl, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl.

25

[00010] The present invention relates to a selective androgen receptor modulator compound having in-vivo androgenic and anabolic activity of a nonsteroidal ligand for

the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula II:



5

wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;

Z is a hydrogen bond acceptor, NO<sub>2</sub>, CN, COR, CONHR;


Y is a lipid soluble group, I, CF<sub>3</sub>, Br, Cl, SnR<sub>3</sub>;

R is an alkyl or aryl group or OH; and

10

Q is acetamido-, trifluoroacetamido-, alkylamines, ether, alkyl, N-sulfonyl, O-sulfonyl, alkylsulfonyl, carbonyl, or a ketone.

[00011] The present invention also relates to a selective androgen receptor modulator compound having in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor the, selective androgen receptor modulator compound represented by the structure of formula III:



where

X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;

Z is NO<sub>2</sub>, CN, COR, or CONHR;

Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;


20

R is an alkyl, or aryl group or OH; and

Q is acetamido or trifluoroacetamido.

[00012] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a

nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula IV:




IV

5

[00013] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula V:


10



V


[00014] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula VI:

15



VI

{00015} The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula VII:



10 {00016} The present invention also relates to a method of binding a selective androgen receptor modulator compound to an androgen receptor, which includes contacting the androgen receptor with the selective androgen receptor modulator compound under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor. In one embodiment the compound is Compound I. In another  
15 embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

20

{00017} Another aspect of the present invention relates to a method of modulating spermatogenesis in a subject, which includes contacting an androgen receptor of the subject with a selective androgen receptor modulator compound under conditions effective to increase or decrease sperm production. In one embodiment the compound is  
25 Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another

embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00018] The present invention also relates to a method of hormone therapy, comprising  
5 contacting an androgen receptor of a subject with a selective androgen receptor  
modulator compound under conditions effective to bind the selective androgen receptor  
modulator compound to the androgen receptor and effect a change in an androgen-  
dependent condition. In one embodiment the compound is Compound I. In another  
embodiment the compound is Compound II. In another embodiment the compound is  
10 Compound III. In another embodiment the compound is Compound IV. In another  
embodiment the compound is Compound V. In another embodiment the compound is  
Compound VI. In another embodiment the compound is Compound VII. In another  
embodiment the compound is Compound VIII.

15 [00019] The present invention also relates to a method of treating a subject having a  
hormone related condition which comprises contacting an androgen receptor of said  
subject with a selective androgen receptor modulator compound under conditions  
effective to bind the selective androgen receptor modulator compound to the androgen  
receptor and effect a change in an androgen-dependent condition. In one embodiment,  
20 the selective androgen receptor modulator compound is selective for androgen or  
testosterone receptor. The present invention also relates to a method of oral  
administration of the selective androgen receptor modulator compound.

[00020] The present invention also relates to a method of treating a subject having a  
25 chronic muscle wasting disease state which comprises contacting an androgen receptor  
of said subject with a selective androgen receptor modulator compound as described  
herein under conditions effective to bind the selective androgen receptor modulator  
compound to the androgen receptor and effect a change in an androgen-dependent  
condition. In one embodiment, the selective androgen receptor modulator compound is  
30 selective for androgen or testosterone receptor. The present invention also relates to a  
method of oral administration of the selective androgen receptor modulator compound.  
In one embodiment the compound is Compound I. In another embodiment the




compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00021] As defined herein, the disease state of "chronic muscle wasting" means

[00022] The present invention also relates to a method of treating a subject having prostate cancer which comprises administering to a subject an effective amount of a selective androgen receptor modulator compound. In one embodiment, the selective androgen receptor modulator compound is selective for androgen or testosterone receptor. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00023] The present invention also relates to compositions and a pharmaceutical compositions which comprises a selective androgen receptor modulator alone or in combination with a progestin or estrogen and a suitable carrier, diluent or salt. In one embodiment the composition comprises Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.


[00024] The present invention relates to a non-steroidal agonist compound, the non-steroidal agonist compound represented by the structure of formula VIII:



wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;  
 R<sub>1</sub> is CH<sub>3</sub>, CF<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, or CF<sub>2</sub>CF<sub>3</sub>;  
 T is OH, OR, -NHCOCH<sub>3</sub>, or NHCOR wherein R is a C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl;


5

A is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or heterocyclic ring represented by the structure:



10

B is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or heterocyclic ring represented by the structure:



15

wherein A<sub>1</sub>-A<sub>11</sub> are each C, O, S or N;  
 B<sub>1</sub>-B<sub>11</sub> are each C, O, S or N;  
 Z is NO<sub>2</sub>, CN, COOH COR, or CONHR;  
 Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>; and

20

Q<sub>1</sub> and Q<sub>2</sub> are independently of each other alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHCSO<sub>2</sub>CH<sub>3</sub>, NHCSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR wherein R is a C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl.

25

[00025] The present invention also relates to a composition and pharmaceutical composition comprising the non-steroidal agonist compound alone or in combination with a progestin or estrogen and a suitable carrier, diluent or salt. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

10 [00026] The present invention also relates to a method of binding a non-steroidal agonist compound to an androgen receptor comprising contacting the androgen receptor with the non-steroidal agonist compound under conditions effective to bind the non-steroidal agonist compound to the androgen receptor. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

20 [00027] The present invention also relates to a method of modulating spermatogenesis in a subject comprising contacting an androgen receptor of the subject with a non-steroidal agonist compound under conditions effective to increase or decrease sperm production. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

30 [00028] The present invention also relates to a method of hormone therapy comprising contacting an androgen receptor of a subject with a non-steroidal agonist under conditions effective to bind the non-steroidal agonist compound to the androgen receptor

and effect a change in an androgen-dependent condition. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In  
5 another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00029] The present invention also relates to a method of treating a subject having a hormone related condition which comprises contacting an androgen receptor of said  
10 subject with a non-steroidal agonist compound under conditions effective to bind the non-steroidal agonist compound to the androgen receptor and effect a change in an androgen-dependent condition. In one embodiment, the non-steroidal agonist compound is selective for androgen or testosterone receptor. The present invention also relates to a method of oral administration of the non-steroidal agonist compound. In one  
15 embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is  
20 Compound VIII.

[00030] The present invention also relates to a method of treating a subject having prostate cancer which comprises administering to a subject an effective amount of a non-steroidal agonist compound. In one embodiment, the non-steroidal agonist  
25 compound is selective for androgen or testosterone receptor. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the  
30 compound is Compound VII. In another embodiment the compound is Compound VIII.

[00031] Still another aspect of the present relates to a method of producing a selective

androgen receptor modulator or a non-steroidal AR agonist compound of the present invention. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00032] The present invention further relates to a method of determining the presence of a selective androgen modulator compound and/or a non-steroidal agonist compound of the present invention in a sample. The method comprises the steps of obtaining the sample, and detecting the compound in the sample, thereby determining the presence of the compound in the sample. In one embodiment, the sample is a blood serum, plasma, urine, or saliva sample. In another embodiment, the detection step comprises measuring the absorbance of the compound. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00033] The novel selective androgen receptor modulator compounds and the non-steroidal agonist compounds of the present invention, either alone or as a composition, are useful in males and females for the treatment of a variety of hormone-related conditions, such as hypogonadism, sarcopenia, erythropoiesis, erectile function, lack of libido, osteoporosis and fertility. Further, the selective androgen receptor modulator compounds and the non-steroidal agonist compounds are useful for oral testosterone replacement therapy, treating prostate cancer, imaging prostate cancer, and maintaining sexual desire in women. The agents may be used alone or in combination with a progestin or estrogen. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another

embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

5 [00034] The selective androgen receptor modulator compounds and the non-steroidal agonist compounds of the present invention offer a significant advance over steroidal androgen treatment because the selective androgen receptor modulator compounds and the non-steroidal agonist compounds of the present invention have been shown in-vivo to have a tissue-selective androgenic and anabolic activity of a nonsteroidal ligand for  
10 the androgen receptor. Moreover, the selective androgen receptor modulator compounds and the non-steroidal agonist compounds of the present invention are not accompanied by serious side effects, lability to oxidative metabolism, inconvenient modes of administration, or high costs and still have the advantages of oral bioavailability, lack of cross-reactivity with other steroid receptors, and long biological half-lives. In one  
15 embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is  
20 Compound VIII.

### BRIEF DESCRIPTION OF THE DRAWINGS

The present invention will be understood and appreciated more fully from the  
25 following detailed description taken in conjunction with the appended drawings in which:

**Figure 1:** Androgenic and Anabolic activity of (S)-GTx-007 in rats. Rats were left untreated (intact control), castrated (castrated control), treated with  
30 testosterone propionate (TP), or treated with S-GTx-007, and the body weight gain as well as the weight of androgen-responsive tissues (prostate, seminal vesicles and levator ani muscle) was determined.

**Figure 2:** Androgenic and Anabolic activity of S-GTx-007 in rats. Rats were left untreated (intact control), castrated (castrated control), treated with 0.1, 0.3, 0.5, 0.75 and 1.0 mg/day testosterone propionate (TP), or treated with 0.1, 0.3, 0.5, 0.75 and 1.0 mg/day S-GTx-007, and the weight of androgen-responsive tissues (prostate, seminal vesicles and levator ani muscle) was determined.

**Figure 3:** Androgenic and Anabolic activity of S-GTx-014 in rats. Rats were left untreated (intact control), castrated (castrated control), treated with 0.1, 0.3, 0.5, 0.75 and 1.0 mg/day testosterone propionate (TP), or treated with 0.1, 0.3, 0.5, 0.75 and 1.0 mg/day S-GTx-014, and the weight of androgen-responsive tissues (prostate, seminal vesicles and levator ani muscle) was determined.

**Figure 4:** Average plasma concentration-time profiles of S-GTx-007 in beagle dogs after IV administration at 3 and 10 mg/kg.

**Figure 5:** Average plasma concentration-time profiles of S-GTx-007 in beagle dogs after PO administration as solution at 10 mg/kg.

**Fig 6:** Average plasma concentration-time profiles of S-GTx-007 in beagle dogs after IV administration as capsules at mg/kg.

**Figure 7:** Effects of GTx-014 and GTx-007 on LH Levels.

**Figure 8:** Effects of GTx-014 and GTx-007 on FSH Levels.

**Figure 9:** Synthesis scheme of GTX-007.

### DETAILED DESCRIPTION OF THE INVENTION

[00035] This invention provides a novel class of androgen receptor targeting agents (ARTA). The agents define a new subclass of compounds which are tissue-selective androgen receptor modulators (SARM) which are useful for oral testosterone replacement therapy, male contraception, maintaining sexual desire in women, treating prostate cancer and imaging prostate cancer. These agents have an unexpected tissue-selective in-vivo activity for an androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor. These agents may be active alone or in combination with progestins or estrogens. The invention further provides a novel class of non-steroidal agonist compounds. The invention further provides compositions containing the selective androgen modulator compounds or the non-steroidal agonist compounds and methods of binding an androgen receptor, modulating spermatogenesis, treating and imaging prostate cancer, and providing hormonal therapy for androgen-dependent conditions.


[00036] The compounds described herein, define a new class of selective androgen receptor modulators (SARMS) that demonstrate potent anabolic effects (e.g., muscle growth) with less androgenic activity (e.g., prostatic growth). This new class of drugs has several advantages over non-selective androgens, including potential therapeutic applications in males and females for modulation of fertility, erythropoiesis, osteoporosis, sexual libido and in men with or at high risk for prostate cancer. In one embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

[00037] Further, in one embodiment the compounds have tissue specific pharmacologic activity. As demonstrated in Figure 7 and 8, GTx-007 does not suppress LH levels at doses that are capable of eliciting maximal stimulation of levator ani muscle growth and



does not suppress FSH levels at doses that are capable of eliciting maximal stimulation of levator ani muscle growth.

[00038] The present invention relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor selective androgen receptor modulator compound represented by the structure of formula I:




wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;


Z is NO<sub>2</sub>, CN, COR, COOH or CONHR;

Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;


Q is alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR wherein R is an aryl, C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl; or Q together with the benzene ring to which it is attached is a fused ring system represented by structure A, B or C:



A



B




C

$R_1$  is  $\text{CH}_3$ ,  $\text{CF}_3$ ,  $\text{CH}_2\text{CH}_3$ , or  $\text{CF}_2\text{CF}_3$ ; and

T is OH, OR,  $-\text{NHCOCH}_3$ , or  $\text{NHCOR}$  wherein R is a  $\text{C}_1$ - $\text{C}_4$  alkyl, a  $\text{C}_1$ - $\text{C}_4$  haloalkyl, phenyl, halo, alkenyl or hydroxyl.

[00039] In one embodiment, Q is in the para position of the benzene ring to which it is attached. In another embodiment, Q is in the para position and X is O. In another embodiment, Q is in the para position and is alkyl, halogen,  $\text{NR}_2$ ,  $\text{NHCOCH}_3$ ,  $\text{NHCOCF}_3$ ,  $\text{NHCOR}$ ,  $\text{NHCONHR}$ ,  $\text{NHCOOR}$ ,  $\text{OCONHR}$ ,  $\text{CONHR}$ ,  $\text{NHCSCH}_3$ ,  $\text{NHCSCF}_3$ ,  $\text{NHCSR}$ ,  $\text{NHSO}_2\text{CH}_3$ ,  $\text{NHSO}_2\text{R}$ , OR, COR, OCOR,  $\text{OSO}_2\text{R}$ ,  $\text{SO}_2\text{R}$  or SR wherein R is a, aryl,  $\text{C}_1$ - $\text{C}_4$  alkyl, a  $\text{C}_1$ - $\text{C}_4$  haloalkyl, phenyl, halo, alkenyl or hydroxyl.

[00040] The present invention relates to a selective androgen receptor modulator compound having in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula II:



20

wherein X is a O,  $\text{CH}_2$ , NH, Se, PR, or NR;


Z is a hydrogen bond acceptor,  $\text{NO}_2$ , CN, COR, CONHR;

Y is a lipid soluble group, I,  $\text{CF}_3$ , Br, Cl,  $\text{SnR}_3$ ;

R is an alkyl, or aryl group or OH; and

25 Q is acetamido-, trifluoroacetamido-, alkylamines, ether, alkyl, N-sulfonyl, O-sulfonyl, alkylsulfonyl, carbonyl, or a ketone.


[00041] The present invention also relates to a selective androgen receptor modulator compound having in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor the, selective androgen receptor modulator compound represented by the structure of formula III:



5                    where        X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;  
                                       Z is NO<sub>2</sub>, CN, COR, or CONHR;  
                                       Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;  
                                       R is an alkyl or aryl group or OH; and  
                                       Q is acetamido or trifluoroacetamido.


10

[00042] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula IV:




IV

20    [00043] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula V:




V

[00044] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a  
5 nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula VI:




VI

[00045] The present invention also relates to a selective androgen receptor modulator compound having tissue-selective in-vivo androgenic and anabolic activity of a  
10 nonsteroidal ligand for the androgen receptor, the selective androgen receptor modulator compound represented by the structure of formula VII:




VII

[00046] The present invention relates to a non-steroidal agonist compound having the  
15 formula (Compound VIII):




VIII

- 5        wherein    X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;  
                   R<sub>1</sub> is CH<sub>3</sub>, CF<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, or CF<sub>2</sub>CF<sub>3</sub>;  
                   T is OH, OR, -NHCOCH<sub>3</sub>, or NHCOR wherein R is a C<sub>1</sub>-C<sub>4</sub>  
                   alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl;  
                   A is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or  
 10        heterocyclic ring represented by the structure:



B is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or  
 heterocyclic ring represented by the structure:

15



wherein       $A_1$ -  $A_{11}$  are each C, O, S or N;  
                  $B_1$ -  $B_{11}$  are each C, O, S or N;  
5                Z is  $\text{NO}_2$ , CN, COR, COOH, or CONHR;  
                 Y is I,  $\text{CF}_3$ , Br, Cl, or  $\text{SnR}_3$ ; and  
                  $Q_1$  and  $Q_2$  are independently of each other alkyl, halogen,  $\text{NR}_2$ ,  
                  $\text{NHCOCH}_3$ ,  $\text{NHCOCF}_3$ ,  $\text{NHCOR}$ ,  $\text{NHCONHR}$ ,  $\text{NHCOOR}$ ,  $\text{OCONHR}$ ,  $\text{CONHR}$ ,  
                  $\text{NHCSCH}_3$ ,  $\text{NHCSCF}_3$ ,  $\text{NHCSR}$ ,  $\text{NHSO}_2\text{CH}_3$ ,  $\text{NHSO}_2\text{R}$ , OR, COR, OCOR,  
10                 $\text{OSO}_2\text{R}$ ,  $\text{SO}_2\text{R}$  or SR wherein R is a  $\text{C}_1$ - $\text{C}_4$  alkyl, a  $\text{C}_1$ - $\text{C}_4$  haloalkyl, phenyl, halo,  
                 alkenyl or hydroxyl.

[00047] The substituents Z and Y can be in any position of the five or 6 membered ring carrying these substituents (hereinafter "A ring"). Similarly, the substituent Q can be in  
15 any position of the five or 6 membered ring carrying this substituent (hereinafter "B ring"). It is understood that when any of the ring members  $A_1$ -  $A_{11}$  or  $B_1$ -  $B_{11}$  are O or S, then these ring members are unsubstituted. It is further understood that when any of the ring members  $A_1$ -  $A_{11}$  or  $B_1$ -  $B_{11}$  are O or S, then the dotted line between said ring members and other ring members represents a single bond.

20

[00048] In one embodiment, the A ring includes any type of saturated or unsaturated carbocyclic ring. In one embodiment, the A ring is a 6 membered saturated carbocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In one embodiment, the A ring is a 5 membered  
25 saturated carbocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 6 membered carbocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 5  
30 membered carbocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove.

[00049] In another embodiment, the A ring includes any type of saturated, unsaturated or aromatic heterocyclic ring. In another embodiment, the A ring is a 6 membered saturated heterocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 5 membered saturated heterocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 6 membered heterocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 5 membered heterocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 6 membered heteroaromatic ring which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the A ring is a 5 membered heteroaromatic ring which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove.

[00050] Similarly, the B ring includes any type of saturated or unsaturated carbocyclic ring. In one embodiment, the B ring is a 6 membered saturated carbocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In one embodiment, the B ring is a 5 membered saturated carbocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the B ring is a 6 membered carbocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the B ring is a 5 membered carbocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove.

[00051] In another embodiment, the B ring includes any type of saturated, unsaturated or aromatic heterocyclic ring. In another embodiment, the B ring is a 6 membered saturated heterocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another  
5 embodiment, the B ring is a 5 membered saturated heterocyclic ring, which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the B ring is a 6 membered heterocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another  
10 embodiment, the B ring is a 5 membered heterocyclic ring containing one or more double bonds, which ring may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another embodiment, the B ring is a 6 membered heteroaromatic ring which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove. In another  
15 embodiment, the B ring is a 5 membered heteroaromatic ring which may be unsubstituted, monosubstituted or polysubstituted by any of the substituents described hereinabove.

Nonlimiting examples of suitable A rings and/or B rings are carbocyclic rings such as  
20 cyclopentane, cyclopentene, cyclohexane, and cyclohexene rings, and heterocyclic rings such as pyran, dihydropyran, tetrahydropyran, pyrrole, dihydropyrrole, tetrahydropyrrole, pyrazine, dihydropyrazine, tetrahydropyrazine, pyrimidine, dihydropyrimidine, tetrahydropyrimidine, pyrazol, dihydropyrazol, tetrahydropyrazol, piperidine, piperazine, pyridine, dihydropyridine, tetrahydropyridine, morpholine,  
25 thiomorpholine, furan, dihydrofuran, tetrahydrofuran, thiophene, dihydrothiophene, tetrahydrothiophene, thiazole, imidazole, isoxazole, and the like.

[00052] As used herein, receptors for extracellular signaling molecules are collectively referred to as "cell signaling receptors". Many cell signaling receptors are  
30 transmembrane proteins on a cell surface; when they bind an extracellular signaling molecule (i.e., a ligand), they become activated so as to generate a cascade of intracellular signals that alter the behavior of the cell. In contrast, in some cases, the



receptors are inside the cell and the signaling ligand has to enter the cell to activate them; these signaling molecules therefore must be sufficiently small and hydrophobic to diffuse across the plasma membrane of the cell. As used herein, these receptors are collectively referred to as "intracellular cell signaling receptors".

5

[00053] Steroid hormones are one example of small hydrophobic molecules that diffuse directly across or are transported across the plasma membrane of target cells and bind to intracellular cell signaling receptors. These receptors are structurally related and constitute the intracellular receptor superfamily (or steroid-hormone receptor  
10 superfamily). Steroid hormone receptors include progesterone receptors, estrogen receptors, androgen receptors, glucocorticoid receptors, and mineralocorticoid, and numerous orphan receptors. The present invention is particularly directed to androgen receptors and all of its isoforms.

15 [00054] In addition to ligand binding to the receptors, the receptors can be blocked to prevent ligand binding. When a substance binds to a receptor, the three-dimensional structure of the substance fits into a space created by the three-dimensional structure of the receptor in a ball and socket configuration.

20 [00055] The better the ball fits into the socket, the more tightly it is held. This phenomenon is called affinity. If the affinity of a substance is sufficiently high, it will compete with the hormone and bind the binding site more frequently. The binding of the ligand may also lead to tissue-selective recruitment of other important proteins to transduce the signal. These proteins are known as coactivators and corepressor,  
25 participate in signal transduction, and may be selectively induced or inhibited by ligand binding. Once bound, signals may be sent through the receptor into the cells, causing the cell to respond in some fashion. This is called activation. On activation, the activated receptor then directly regulates the transcription of specific genes. But the substance and the receptor may have certain attributes, other than affinity, that activate the cell.  
30 Chemical bonds between atoms of the substance and the atoms of the receptors may form. In some cases, this leads to a change in the configuration of the receptor, which is enough to begin the activation process (called signal transduction). As a result,

substances can be made which bind receptors and activate them (called receptor agonists) or inactivate them (called receptor antagonists).

[00056] The present invention is directed to selective androgen receptor modulator compounds which are agonist compounds, and are, therefore, useful in binding to and activating steroidal hormone receptors. The compounds are non-steroidal. Preferably, the agonist compound of the present invention is an agonist that binds the androgen receptor. Preferably, the compound has high affinity for the androgen receptor. The compound may bind either reversibly or irreversibly to the androgen receptor. The compound of the present invention may contain a functional group (affinity label) that allows alkylation of the androgen receptor (i.e. covalent bond formation). Thus, in this case, the compound binds irreversibly to the receptor and, accordingly, cannot be displaced by a steroid, such as the endogenous ligands dihydrotestosterone and testosterone. It is preferable, however, for the compounds of the present invention to reversibly bind the androgen receptor.

[00057] According to one aspect of the present invention, a method is provided for binding the selective androgen receptor modulator compounds of the present invention to an androgen receptor by contacting the receptor with a selective androgen receptor modulator compound under conditions effective to cause the selective androgen receptor modulator compound to bind the androgen receptor. The binding of the selective androgen receptor modulator compounds to the androgen receptor enables the compounds of the present invention to be useful in males and in females in a number of hormone therapies. The agonist compounds bind to and activate the androgen receptor. Binding of the agonist compound is either reversible or irreversible, preferably reversible.

[00058] According to one aspect of the present invention, a method is provided for modulating spermatogenesis by contacting an androgen receptor of a patient with a selective androgen receptor modulator compound under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor and increase or decrease sperm production.

[00059] According to another aspect of the present invention, a method is provided for hormonal therapy in a patient (i.e., suffering from an androgen- dependent condition) which includes contacting an androgen receptor of a patient with a selective androgen  
5 receptor modulator compound under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor and effect a change in an androgen-dependent condition. Androgen-dependent conditions that may be treated according to the present invention include those conditions associated with aging, such as hypogonadism, sarcopenia, erythropoiesis, osteoporosis, and any other conditions  
10 later determined to be dependent upon low androgen (e.g., testosterone) levels. In one embodiment, the selective androgen receptor modulator compound is administered alone. In another embodiment, the selective androgen receptor modulator compound is administered in combination with progestin. In yet another embodiment, the selective androgen receptor modulator compound is administered in combination with estrogen.

15

[00060] According to another aspect of the present invention, a method is provided for treating a subject having prostate cancer. The method comprises administering to a subject an effective amount of a selective androgen receptor modulator compound. In one embodiment, the selective androgen receptor modulator compound is selective for  
20 androgen or testosterone receptor.

[00061] The present invention also relates to a method of treating a subject having a chronic muscle wasting disease state which comprises contacting an androgen receptor of said subject with a selective androgen receptor modulator compound under conditions  
25 effective to bind the selective androgen receptor modulator compound to the androgen receptor and effect a change in an androgen-dependent condition. In one embodiment, the selective androgen receptor modulator compound is selective for androgen or testosterone receptor. The present invention also relates to a method of oral administration of the selective androgen receptor modulator compound. In one  
30 embodiment the compound is Compound I. In another embodiment the compound is Compound II. In another embodiment the compound is Compound III. In another embodiment the compound is Compound IV. In another embodiment the compound is

Compound V. In another embodiment the compound is Compound VI. In another embodiment the compound is Compound VII. In another embodiment the compound is Compound VIII.

5

[00062] According to one aspect of the present invention, a method is provided for binding the non-steroidal agonist compounds of the present invention to an androgen receptor by contacting the receptor with a non-steroidal agonist compound under conditions effective to cause the non-steroidal agonist compound to bind the androgen  
10 receptor. The binding of the non-steroidal agonist compounds to the androgen receptor enables the compounds of the present invention to be useful in males and in females in a number of hormone therapies. The agonist compounds bind to and activate the androgen receptor. Binding of the agonist compound is either reversible or irreversible, preferably reversible.

15

[00063] According to one aspect of the present invention, a method is provided for modulating spermatogenesis by contacting an androgen receptor of a patient with a non-steroidal agonist compound under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor and increase or decrease sperm  
20 production.

[00064] According to another aspect of the present invention, a method is provided for hormonal therapy in a patient (i.e., one suffering from an androgen-dependent condition) which includes contacting an androgen receptor of a patient with a non-steroidal agonist compound under conditions effective to bind the non-steroidal agonist  
25 compound to the androgen receptor and effect a change in an androgen-dependent condition. Androgen-dependent conditions that may be treated according to the present invention include those conditions associated with aging, such as hypogonadism, sarcopenia, erythropoiesis, osteoporosis, lack of sexual libido and any other conditions  
30 later determined to be dependent upon low androgen (e.g., testosterone) levels. In one embodiment, the non-steroidal agonist compound is administered alone. In another embodiment, the non-steroidal agonist compound is administered in combination with

progesterin. In yet another embodiment, the non-steroidal agonist compound is administered in combination with estrogen.

5 [00065] According to another aspect of the present invention, a method is provided for treating a subject having prostate cancer. The method comprises administering to a subject an effective amount of a non-steroidal agonist compound. In one embodiment, the non-steroidal agonist compound is selective for androgen or testosterone receptor.

10 [00066] The compounds of the present invention have an asymmetric center and can be the R or S isomer, or a mixture of both. In one embodiment, the compounds are racemic mixtures of the R and S enantiomers. In another embodiment, the compounds are substantially pure R enantiomers. In another embodiment, the compounds are substantially pure S enantiomers. "Substantially pure" is defined herein as greater than about 95% preponderance of one isomer. Where the above-described processes for the  
15 preparation of the compounds of use in the invention give rise to mixtures of stereoisomers, these isomers may be separated by conventional techniques, such as preparative chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution

20

[00067] As used herein, "pharmaceutical composition" means therapeutically effective amounts of the SARM or the non-steroidal agonist compound of the present invention, together with suitable diluents, preservatives, solubilizers, emulsifiers, adjuvant and/or carriers. A "therapeutically effective amount" as used herein refers to that amount which  
25 provides a therapeutic effect for a given condition and administration regimen. Such compositions are liquids or Lyophilized or otherwise dried formulations and include diluents of various buffer content (e.g., Tris-HCl, acetate, phosphate), pH and ionic strength, additives such as albumin or gelatin to prevent adsorption to surfaces, detergents (e.g., Tween 20, Tween 80, Pluronic F68, bile acid salts), solubilizing agents  
30 (e.g., glycerol, polyethylene glycerol), anti-oxidants (e.g., ascorbic acid, sodium metabisulfite), preservatives (e.g., Thimerosal, benzyl alcohol, parabens), bulking substances or tonicity modifiers (e.g., lactose, mannitol), covalent attachment of

polymers such as polyethylene glycol to the protein, complexation with metal ions, or incorporation of the material into or onto particulate preparations of polymeric compounds such as polylactic acid, polglycolic acid, hydrogels, etc, or onto liposomes, microemulsions, micelles, unilamellar or multilamellar vesicles, erythrocyte ghosts, or spheroplasts. Such compositions will influence the physical state, solubility, stability, rate of in vivo release, and rate of in vivo clearance. Controlled or sustained release compositions include formulation in lipophilic depots (e.g., fatty acids, waxes, oils).

[00068] Also comprehended by the invention are particulate compositions coated with polymers (e.g., poloxamers or poloxamines). Other embodiments of the compositions of the invention incorporate particulate forms, protective coatings, protease inhibitors or permeation enhancers for various routes of administration, including parenteral, pulmonary, nasal and oral. In one embodiment the pharmaceutical composition is administered parenterally, paracancerally, transmucosally, transdermally, intramuscularly, intravenously, intradermally, subcutaneously, intraperitoneally, intraventricularly, intracranially and intratumorally.

[00069] Further, as used herein "pharmaceutically acceptable carriers" are well known to those skilled in the art and include, but are not limited to, 0.01-0.1M and preferably 0.05M phosphate buffer or 0.8% saline. Additionally, such pharmaceutically acceptable carriers may be aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media.

[00070] Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers such as those based on Ringer's dextrose, and the like. Preservatives and other additives may also be present, such as, for example, antimicrobials, antioxidants, collating agents, inert gases and the like.

[00071] Controlled or sustained release compositions include formulation in lipophilic depots (e.g. fatty acids, waxes, oils). Also comprehended by the invention are particulate compositions coated with polymers (e.g. poloxamers or poloxamines) and the compound  
5 coupled to antibodies directed against tissue-specific receptors, ligands or antigens or coupled to ligands of tissue-specific receptors.

[00072] Other embodiments of the compositions of the invention incorporate particulate forms, protective coatings, protease inhibitors or permeation enhancers for  
10 various routes of administration, including parenteral, pulmonary, nasal and oral.

[00073] Compounds modified by the covalent attachment of water-soluble polymers such as polyethylene glycol, copolymers of polyethylene glycol and polypropylene glycol, carboxymethyl cellulose, dextran, polyvinyl alcohol, polyvinylpyrrolidone or  
15 polyproline are known to exhibit substantially longer half-lives in blood following intravenous injection than do the corresponding unmodified compounds (Abuchowski et al., 1981; Newmark et al., 1982; and Katre et al., 1987). Such modifications may also increase the compound's solubility in aqueous solution, eliminate aggregation, enhance the physical and chemical stability of the compound, and greatly reduce the  
20 immunogenicity and reactivity of the compound. As a result, the desired in vivo biological activity may be achieved by the administration of such polymer-compound adducts less frequently or in lower doses than with the unmodified compound.

[00074] In yet another embodiment, the pharmaceutical composition can be delivered  
25 in a controlled release system. For example, the agent may be administered using intravenous infusion, an implantable osmotic pump, a transdermal patch, liposomes, or other modes of administration. In one embodiment, a pump may be used (see Langer, supra; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)). In another embodiment,  
30 polymeric materials can be used. In yet another embodiment, a controlled release system can be placed in proximity to the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled

Release, *supra*, vol. 2, pp. 115-138 (1984). Preferably, a controlled release device is introduced into a subject in proximity to the site of inappropriate immune activation or a tumor. Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

5

[00075] The pharmaceutical preparation can comprise the selective androgen receptor modulator alone, or can further include a pharmaceutically acceptable carrier, and can be in solid or liquid form such as tablets, powders, capsules, pellets, solutions, suspensions, elixirs, emulsions, gels, creams, or suppositories, including rectal and urethral  
10 suppositories. Pharmaceutically acceptable carriers include gums, starches, sugars, cellulosic materials, and mixtures thereof. The pharmaceutical preparation containing the selective androgen receptor modulator can be administered to a subject by, for example, subcutaneous implantation of a pellet; in a further embodiment, the pellet provides for controlled release of selective androgen receptor modulator over a period of  
15 time. The preparation can also be administered by intravenous, intraarterial, or intramuscular injection of a liquid preparation, oral administration of a liquid or solid preparation, or by topical application. Administration can also be accomplished by use of a rectal suppository or a urethral suppository.

20 [00076] The pharmaceutical preparations of the invention can be prepared by known dissolving, mixing, granulating, or tablet-forming processes. For oral administration, the selective androgen receptor modulators or their physiologically tolerated derivatives such as salts, esters, N-oxides, and the like are mixed with additives customary for this purpose, such as vehicles, stabilizers, or inert diluents, and converted by customary  
25 methods into a suitable form for administration, such as tablets, coated tablets, hard or soft gelatin capsules, aqueous, alcoholic or oily solutions. Examples of suitable inert vehicles are conventional tablet bases such as lactose, sucrose, or cornstarch in combination with binders like acacia, cornstarch, gelatin, or with disintegrating agents such as cornstarch, potato starch, alginic acid, or with a lubricant like stearic acid or  
30 magnesium stearate.

[00077] Examples of suitable oily vehicles or solvents are vegetable or animal oils



such as sunflower oil or fish-liver oil. Preparations can be effected both as dry and as wet granules. For parenteral administration (subcutaneous, intravenous, intraarterial, or intramuscular injection), the SARM agents or the non-steroidal agonist agents or their physiologically tolerated derivatives such as salts, esters, N-oxides, and the like are  
5 converted into a solution, suspension, or emulsion, if desired with the substances customary and suitable for this purpose, for example, solubilizers or other auxiliaries. Examples are: sterile liquids such as water and oils, with or without the addition of a surfactant and other pharmaceutically acceptable adjuvants. Illustrative oils are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, or  
10 mineral oil. In general, water, saline, aqueous dextrose and related sugar solutions, and glycols such as propylene glycols or polyethylene glycol are preferred liquid carriers, particularly for injectable solutions.

[00078] The preparation of pharmaceutical compositions which contain an active  
15 component is well understood in the art. Typically, such compositions are prepared as aerosols of the polypeptide delivered to the nasopharynx or as injectables, either as liquid solutions or suspensions, however, solid forms suitable for solution in, or suspension in, liquid prior to injection can also be prepared. The preparation can also be emulsified. The active therapeutic ingredient is often mixed with excipients that are  
20 pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof.

[00079] In addition, if desired, the composition can contain minor amounts of auxiliary  
25 substances such as wetting or emulsifying agents, pH buffering agents, which enhance the effectiveness of the active ingredient.

[00080] An active component can be formulated into the composition as neutralized  
pharmaceutically acceptable salt forms. Pharmaceutically acceptable salts include the  
30 acid addition salts (formed with the free amino groups of the polypeptide or antibody molecule), which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like.

Salts formed from the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

5

[00081] For topical administration to body surfaces using, for example, creams, gels, drops, and the like, the SARM agents or the non-agonist steroidal compounds or their physiologically tolerated derivatives such as salts, esters, N-oxides, and the like are prepared and applied as solutions, suspensions, or emulsions in a physiologically acceptable diluent with or without a pharmaceutical carrier.

10

[00082] In another embodiment, the active compound can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*).

15

[00083] For use in medicine, the salts of the SARM or the non-steroidal agonist compounds will be pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds according to the invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts, which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, methanesulphonic acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, oxalic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid.

20

25

[00084] The present invention further relates to a method of determining the presence of a selective androgen modulator compound and/or a non-steroidal agonist compound of the present invention in a sample. The method comprises the steps of obtaining the sample, and detecting the compound in the sample, thereby determining the presence of the compound in the sample.

30

[00085] In one embodiment, the sample is a blood serum sample. In another embodiment, the sample is a plasma sample. In another embodiment, the sample is a urine sample. In another embodiment, the sample is a saliva sample. In another  
5 embodiment, the sample is any other tissue sample.

[00086] In one embodiment, the detection step comprises measuring the absorbance of the compound at a predetermined wavelength. For example, the compounds of the present invention absorb in the ultraviolet region of the spectrum, with an absorbency  
10 peak at 270 nm. Thus, in one embodiment of the present invention, the compound is detected by monitoring the UV absorbance of the sample at 270 nm. It should be noted that the present invention is not limited to UV absorption, and that any other spectrometric methods of identification are applicable. For example, compounds can be detected by measuring their infra-red or visible absorbance.

15

[00087] In another embodiment, the present invention further provides a method of determining the concentration of a selective androgen receptor modulator compound and/or a non-steroidal agonist compound of the present invention in a sample. The method comprises the steps of obtaining a sample; determining the level of the  
20 compound in the sample, and calculating the concentration of the compound in the sample by comparing the level with a standard sample containing a known concentration of the compound. Calibration curves of known concentrations of the compound in the sample, can be obtained, and the concentration of the compound in the test sample is calculated therefrom. By "level" it is meant the absorption level of the compound at the  
25 measured wavelength.

[00088] In another embodiment, the compound is detected in the sample by contacting the sample with a binding protein which specifically binds to the compound, and determining the amount of binding protein bound to the compound. The concentration  
30 of the compound can be determined by measuring the amount of binding protein bound to the compound, and comparing that amount to a standard sample containing a known concentration of the compound – binding protein complex.

[00089] Protein levels can be determined according to standard techniques, as described in Sambrook et al. Briefly, a sample obtained from a subject is contacted with a binding protein which specifically binds to a specific compound of the present invention, and the amount of complex formed between the binding protein and the compound is determined. In one embodiment, the binding protein is an antibody which specifically binds to one or more compounds of the present invention. In another embodiment, the binding protein has a detectable label bound thereto, and the complex between the binding protein-label compound is determined by visualizing the complex.

10

[00090] As defined herein, "contacting" means that the binding protein is introduced into the sample in a test tube, flask, tissue culture, chip, array, plate, microplate, capillary, or the like, and incubated at a temperature and time sufficient to permit the binding component to bind to a cell or a fraction thereof or plasma/serum or a fraction thereof containing the target. Methods for contacting the samples with the binding proteins, or other specific binding components are known to those skilled in the art and may be selected depending on the type of assay protocol to be run. Incubation methods are also standard and are known to those skilled in the art.

[00091] "Visualizing" the complex may be carried out by any means known in the art, including, but not limited to, ELISA, radioimmunoassay, flow cytometry, dot blots, western immunoblotting combined with gel electrophoresis, immunohistochemistry at light and electron pe levels, HPLC and mass spectrometry.

[00092] Either monoclonal or polyclonal antibodies (as well as any recombinant antibodies) specific for the selective androgen modulator compounds or the non-steroidal agonist compounds of the present invention can be used in the various immunoassays. The antibodies may be detectably labeled, utilizing conventional labeling techniques well-known to the art. As used herein, the term "label" refers to a molecule, which may be conjugated or otherwise attached (i.e., covalently or non-covalently) to a binding protein as defined herein. Labels are known to those skilled in the art. Thus, the antibodies may be labeled with radioactive isotopes, non-radioactive

30

isotopic labels, fluorescent labels, enzyme labels, chemiluminescent labels, bioluminescent labels, free radical labels, or bacteriophage labels, using techniques known in the art. Examples of radioisotopic labels are <sup>3</sup>H, <sup>125</sup>I, <sup>131</sup>I, <sup>35</sup>S, <sup>14</sup>C, etc. Examples of non-radioactive isotopic labels are <sup>55</sup>Mn, <sup>56</sup>Fe, etc. Examples of fluorescence labels are fluorescent labels which are directly labeled with the preferred fluorescence label, or fluorescent labels which are indirectly labeled with the preferred fluorescence label. In the last case, the preferred fluorescence label is conjugated to a secondary antibody, which is directed against the first antibody, such as an anti species Ig antibody. Typical fluorescent labels include, but are not limited to a fluorescein label, an isothiocyanate label, a rhodamine label, a phycoerythrin label, etc., for example fluorescein isothiocyanate (FITC, International Biological Supplies, Melbourne, FL), rhodamine, phycoerythrin (P.E., Coulter Corp., Hialeah, FL), phycocyanin, allophycocyanin, phycoerythrin-cyanin dye 5 (PECy5, Coulter), label, a phycocyanin label, an allophycocyanin label, an O-phthaldehyde label, a fluorescamine and Texas Red.

[00093] Examples of enzyme labels include alkaline phosphatase, beta-galactosidase, glucose-6-phosphate dehydrogenase, maleate dehydrogenase, and peroxidase. Two principal types of enzyme immunoassay are the enzyme-linked immunosorbent assay (ELISA), and the homogeneous enzyme immunoassay, also known as enzyme-multiplied immunoassay (EMIT, Syva Corporation, Palo Alto, CA). In the ELISA system, separation may be achieved, for example, by the use of antibodies coupled to a solid phase. The EMIT system depends on deactivation of the enzyme in the tracer-antibody complex; the activity can thus be measured without the need for a separation step.

[00094] Particularly suitable labels include those, which permit analysis by flow cytometry, e.g., fluorochromes. Other suitable detectable labels include those useful in colorimetric enzyme systems, e. g., horseradish peroxidase (HRP) and alkaline phosphatase (AP). Other proximal enzyme systems are known to those of skill in the art, including hexokinase in conjunction with glucose-6-phosphate dehydrogenase.

[00095] Additionally, chemiluminescent compounds may be used as labels. Chemiluminescent labels, such as green fluorescent proteins, blue fluorescent proteins, and variants thereof are known. Also bioluminescence or chemiluminescence can be detected using, respectively, NAD oxidoreductase with luciferase and substrates NADH and FNIN or peroxidase with luminol and substrate peroxide. Typical chemiluminescent compounds include luminol, isoluminol, aromatic acridinium esters, imidazoles, acridinium salts, and oxalate esters. Similarly, bioluminescent compounds may be utilized for labelling, the bioluminescent compounds including luciferin, luciferase, and aequorin. Once labeled, the antibody may be employed to identify and quantify immunologic counterparts (antibody or antigenic polypeptide) utilizing techniques well-known to the art.

[00096] The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way be construed, however, as limiting the broad scope of the invention.

## EXPERIMENTAL DETAILS SECTION

### EXAMPLE 1

#### **Nonsteroidal Ligands with Androgenic and Anabolic Activity**

[00097] The SARM compounds provided herein were designed, synthesized and evaluated for in-vitro and in-vivo pharmacologic activity. The in-vitro androgen receptor binding affinity and ability to maintain androgen dependent tissue growth in castrated animals was studied. Androgenic activity was monitored as the ability of the SARM compounds to maintain and/or stimulate the growth of the prostate and seminal vesicles, as measured by weight. Anabolic activity was monitored as the ability of the SARM compounds to maintain and/or stimulate the growth of the levator ani muscle, as measured by weight.

**Synthetic Procedures of Compounds I-VIII:**

[00098] (2R)-1-Methacryloylpyrrolidin-2-carboxylic Acid (R-129). D-Proline (R-128, 14.93 g, 0.13 mol) was dissolved in 71 mL of 2 N NaOH and cooled in an ice bath; the resulting alkaline solution was diluted with acetone (71 mL). An acetone solution (71 mL) of metacryloyl chloride 127 (13.56 g, 0.13 mol) and 2N NaOH solution (71 mL) were simultaneously added over 40 min to the aqueous solution of D-proline in an ice bath. The pH of the mixture was kept at 10-11°C during the addition of the metacryloyl chloride. After stirring (3 h, room temperature), the mixture was evaporated in vacuo at a temperature at 35-45 °C to remove acetone. The resulting solution was washed with ethyl ether and was acidified to pH 2 with concentrated HCl. The acidic mixture was saturated with NaCl and was extracted with EtOAc (100 mL x 3). The combined extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered through Celite, and evaporated in vacuo to give the crude product as a colorless oil. Recrystallization of the oil from ethyl ether and hexanes afforded 16.2 (68%) of the desired compound as colorless crystals: mp 102-103 °C (lit. [214] mp 102.5-103.5 °C); the NMR spectrum of this compound demonstrated the existence of two rotamers of the title compound. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 5.28 (s) and 5.15 (s) for the first rotamer, 5.15 (s) and 5.03 (s) for the second rotamer (totally 2H for both rotamers, vinyl CH<sub>2</sub>), 4.48-4.44 for the first rotamer, 4.24-4.20 (m) for the second rotamer (totally 1H for both rotamers, CH at the chiral center), 3.57-3.38 (m, 2H, CH<sub>2</sub>), 2.27-2.12 (1H, CH), 1.97-1.72 (m, 6H, CH<sub>2</sub>, CH, Me); <sup>13</sup>C NMR (75 MHz, DMSO-d<sub>6</sub>) δ for major rotamer 173.3, 169.1, 140.9, 116.4, 58.3, 48.7, 28.9, 24.7, 19.5; for minor rotamer 174.0, 170.0, 141.6, 115.2, 60.3, 45.9, 31.0, 22.3, 19.7; IR (KBr) 3437 (OH), 1737 (C=O), 1647 (CO, COOH), 1584, 1508, 1459, 1369, 1348, 1178 cm<sup>-1</sup>; [α]<sub>D</sub><sup>26</sup> +80.8° (c = 1, MeOH); Anal. Calcd. for C<sub>9</sub>H<sub>13</sub>NO<sub>3</sub>: C 59.00, H 7.15, N 7.65. Found: C 59.13, H 7.19, N 7.61.

[00099] (3R,8aR)-3-Bromomethyl-3-methyl-tetrahydro-pyrrolo[2,1-c][1,4]oxazine-1,4-dione (R, R-130). A solution of NBS (23.5g, 0.132 mol) in 100 mL of DMF was added dropwise to a stirred solution of compound R-129 (16.1g, 88 mmol) in 70 mL of DMF under argon at room temperature, and the resulting mixture was stirred 3 days. The solvent was removed in vacuo, and a yellow solid was precipitated.

The solid was suspended in water, stirred overnight at room temperature, filtered, and dried to give 18.6 (81%) (smaller weight when dried ~ 34%) of the title compound as a yellow solid: mp 152-154 °C (lit. [214] mp 107-109 °C for the S-isomer); <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 4.69 (dd, J = 9.6 Hz, J = 6.7 Hz, 1H, CH at the chiral center), 4.02 (d, J = 11.4 Hz, 1H, CHH<sub>a</sub>), 3.86 (d, J = 11.4 Hz, 1H, CHH<sub>b</sub>), 3.53-3.24 (m, 4H, CH<sub>2</sub>), 2.30-2.20 (m, 1H, CH), 2.04-1.72 (m, 3H, CH<sub>2</sub> and CH), 1.56 (s, 2H, Me); <sup>13</sup>C NMR (75 MHz, DMSO-d<sub>6</sub>) δ 167.3, 163.1, 83.9, 57.2, 45.4, 37.8, 29.0, 22.9, 21.6; IR (KBr) 3474, 1745 (C=O), 1687 (C=O), 1448, 1377, 1360, 1308, 1227, 1159, 1062 cm<sup>-1</sup>; [α]<sub>D</sub><sup>26</sup> +124.5° (c = 1.3, chloroform); Anal. Calcd. for C<sub>9</sub>H<sub>12</sub>BrNO<sub>3</sub>: C 41.24, H 4.61, N 5.34. Found: C 41.46, H 4.64, N 5.32.

[000100] **(2R)-3-Bromo-2-hydroxy-2-methylpropanoic Acid (R-131).** A mixture of bromolactone **R-130** (18.5g, 71 mmol) in 300 mL of 24% HBr was heated at reflux for 1 h. The resulting solution was diluted with brine (200 mL), and was extracted with ethyl acetate (100 mL x 4). The combined extracts were washed with saturated NaHCO<sub>3</sub> (100 mL x 4). The aqueous solution was acidified with concentrated HCl to pH = 1, which, in turn, was extracted with ethyl acetate (100 mL x 4). The combined organic solution was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered through Celite, and evaporated in vacuo to dryness. Recrystallization from toluene afforded 10.2 g (86%) of the desired compound as colorless crystals: mp 107-109 °C (lit. [214] mp 109-113 °C for the S-isomer); <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 3.63 (d, J = 10.1 Hz, 1H, CHH<sub>a</sub>), 3.52 (d, J = 10.1 Hz, 1H, CHH<sub>b</sub>), 1.35 (s, 3H, Me); IR (KBr) 3434 (OH), 3300-2500 (COOH), 1730 (C=O), 1449, 1421, 1380, 1292, 1193, 1085 cm<sup>-1</sup>; [α]<sub>D</sub><sup>26</sup> +10.5° (c = 2.6, MeOH); Anal. Calcd. for C<sub>4</sub>H<sub>7</sub>BrO<sub>3</sub>: C 26.25, H 3.86. Found: C 26.28, H 3.75.

[000101] **N-[4-Nitro-3-(trifluoromethyl)phenyl]-(2R)-3-bromo-2-hydroxy-2-methylpropanamide (R-132).** Thionyl chloride (8.6 g, 72 mmol) was added dropwise under argon to a solution of bromoacid **R-131** (11.0 g, 60 mmol) in 70 mL of DMA at -5 to -10 °C. The resulting mixture was stirred for 2 h under the same conditions. A solution of 4-nitro-3-trifluoromethyl-aniline (12.4 g, 60 mmol) in 80 mL of DMA was added dropwise to the above solution, and the resulting mixture was stirred overnight at room temperature. The solvent was removed on Rotavapor using high vacuum oil pump;



the residue was diluted with saturated  $\text{NaHCO}_3$  solution, and extracted with ethyl ether (100 mL x 3). Combined extracts were dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered through Celite, and purified by flash chromatography on silica gel, using methylene chloride as eluent to afford 18.0 g (80%) of the desired compound: mp 98-100 °C ( $R_f$  = 0.2, silica gel,  $\text{CH}_2\text{Cl}_2$ );  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  10.54 (s, 1H, NH), 8.54 (d,  $J$  = 2.1 Hz, 1H, ArH), 8.34 (dd,  $J$  = 9.0 Hz,  $J$  = 2.1 Hz, 1H, ArH), 8.18 (d,  $J$  = 9.0 Hz, 1H, ArH), 6.37 (s, 1H, OH), 3.82 (d,  $J$  = 10.4 Hz, 1H,  $\text{CHH}_a$ ), 3.58 (d,  $J$  = 10.4 Hz, 1H,  $\text{CHH}_b$ ), 1.48 (s, 3H, Me);  $^{13}\text{C}$  NMR (75 MHz,  $\text{DMSO}-d_6$ )  $\delta$  173.6 (C=O), 143.0, 127.2, 123.2, 122.6 (q,  $J$  = 33.0 Hz), 122.0 (q,  $J$  = 271.5 Hz), 118.3 (q,  $J$  = 6.0 Hz), 74.4, 41.4, 24.9; IR (KBr) 3344 (OH), 1680 (C=O), 1599, 1548 (C=C, Ar), 1427, 1363, 1161  $\text{cm}^{-1}$ ; MS (ESI):  $m/z$  370.8 ( $\text{M}$ ) $^+$ ; Anal. Calcd. for  $\text{C}_{11}\text{H}_{10}\text{BrN}_2\text{O}_4$ : C 35.60, H 2.72, N 7.55. Found: C 35.68, H 2.72, N 7.49.

[000102] N-[4-nitro-3-trifluoromethylphenyl]-(2S)-3-[4-(acetylamino)phenoxy]-2-hydroxy-2-methylpropanamide (S-147). The title compound was prepared from compound R-132 (0.37 g, 1.0 mmol), 4-acetamidophenol (0.23 g, 1.5 mmol)  $\text{K}_2\text{CO}_3$  (0.28 g, 2.0 mmol), and 10% of benzyltributylammonium chloride as a phase transfer catalyst in 20 mL of methyl ethyl ketone was heated at reflux overnight under argon. The reaction was followed by TLC, the resulting mixture was filtered through Celite, and concentrated in vacuo to dryness. Purification by flash column chromatography on silica gel (hexanes-ethyl acetate, 3:1) yielded 0.38 g (86%) ( $R_f$  = 0.18 hexanes-ethyl acetate, 3:1) of the desired compound as a light yellow powder: mp 70-74 °C; The solid can be recrystallized from ethyl acetate and hexane);  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  10.62 (s, 1H, NH), 9.75 (s, 1H, NH), 8.56 (d,  $J$  = 1.9 Hz, 1H, ArH), 8.36 (dd,  $J$  = 9.1 Hz,  $J$  = 1.9 Hz, 1H, ArH), 8.18 (d,  $J$  = 9.1 Hz, 1H, ArH), 7.45-7.42 (m, 2H, ArH), 6.85-6.82 (m, 2H, ArH), 6.25 (s, 1H, OH), 4.17 (d,  $J$  = 9.5 Hz, 1H,  $\text{CHH}_a$ ), 3.94 (d,  $J$  = 9.5 Hz, 1H,  $\text{CHH}_b$ ), 1.98 (s, 3H, Me), 1.43 (s, 3H, Me);  $^{13}\text{C}$  NMR (75 MHz,  $\text{DMSO}-d_6$ )  $\delta$  174.6 (C=O), 167.7, 154.2, 143.3, 141.6, 132.8, 127.4, 123.0, 122.7 (q,  $J$  = 33.0 Hz), 122.1 (q,  $J$  = 271.5 Hz), 120.1, 118.3 (q,  $J$  = 6.0 Hz), 114.6, 74.9, 73.8, 23.8, 23.0; IR (KBr) 3364 (OH), 1668 (C=O), 1599, 1512 (C=C, Ar), 1457, 1415, 1351, 1323, 1239, 1150 1046  $\text{cm}^{-1}$ ; MS (ESI):  $m/z$  464.1 ( $\text{M}+\text{Na}$ ) $^+$ ; Anal. Calcd. for  $\text{C}_{19}\text{H}_{18}\text{F}_3\text{N}_3\text{O}_6$ : C 51.71, H 4.11, N 9.52. Found: C 52.33, H 4.40, N 9.01.

[Duane insert]

5

[000103] The in-vitro activity of the SARM compounds, specifically compound VII,  
10 demonstrated high androgen receptor binding affinity ( $K_i = 7.5$  nM). Animal studies  
with the SARM compounds, specifically compound V, demonstrated that it is a potent  
androgenic and anabolic nonsteroidal agent. Four groups of rats were used for these  
studies: (1) intact controls, (2) castrated controls, (3) castrated animals treated with  
testosterone propionate (100  $\mu$ g/day), and (4) castrated animals treated with compound  
15 V (1000  $\mu$ g/day). Testosterone and compound VII were delivered at a constant rate for  
14 days via subcutaneous osmotic pumps.


[000104] The results of these studies are shown in Figure 1. Castration significantly  
reduced the weight of androgenic (e.g., prostate and seminal vesicles) and anabolic (e.g.,  
20 levator ani muscle) tissues, but had little effect on animal body weight (BW). Treatment  
of castrated animals with testosterone propionate or compound VII maintained the  
weight of androgenic tissues to the same degree. Compound VII had similar androgenic  
activity as testosterone propionate (i.e., the prostate and seminal vesicle weights were  
the same), but much greater efficacy as an anabolic agent. Compound VII showed  
25 greater anabolic activity than testosterone propionate at the doses tested (i.e., the levator  
ani muscle maintained the same weight as intact control animals and was greater than  
that observed for testosterone). The experiments presented herein are the first in-vivo  
results which demonstrate tissue-selective androgenic and anabolic activity (i.e.,  
differing androgenic and anabolic potency) of a nonsteroidal ligand for the androgen  
30 receptor.

**EXAMPLE 2****Nonsteroidal Ligands with Androgenic and Anabolic Activity**

5 [000105] The in-vivo efficacy and acute toxicity of four novel nonsteroidal androgens (compounds IV, V, VI and VII) in rats was examined. In-vitro assays established that these compounds bind the androgen receptor with very high affinity. The structures and names of the four compounds are presented below:

10

15



GTx-014	R= F
GTx-015	R=COCH <sub>3</sub>
GTx-016	R=COC <sub>2</sub> H <sub>5</sub>
GTx-017	R=NHCOCH <sub>3</sub>

20

**EXPERIMENTAL METHODS**

[000106] *Materials.* The S-isomers of compounds GTx-014 (compound IV), GTx-015 (compound V), GTx-016 (compound VI) and GTx-007 (compound VII wherein R is NHCOCH<sub>3</sub>) and the R-isomer of GTx-014 were synthesized in accordance with the scheme as set forth in Figure 9. Testosterone propionate (TP), polyethylene glycol 300 (PEG300, reagent grade) and neutral buffered formalin (10% w/v) were purchased from

25

Sigma Chemical Company (St Louis, MO). Alzet osmotic pumps (model 2002) were purchased from Alza Corp. (Palo Alto, CA).

5 [000107] *Animals.* Immature male Sprague-Dawley rats, weighing 90 to 100g, were purchased from Harlan Biosciences (Indianapolis, IN). The animals were maintained on a 12-hour light-dark cycle with food and water available *ad libitum*. The animal protocol was reviewed and approved by the Institutional Laboratory Animal Care and Use Committee.

10 [000108] *Study Design.* Rats were randomly distributed into twenty-nine (29) groups, with 5 animals per group. Treatment groups are described in Table 1. One day prior to the start of drug treatment, animals in groups 2 through 29 were individually removed from the cage, weighed and anesthetized with an intraperitoneal dose of ketamine/xylazine (87/13 mg/kg; approximately 1 mL per kg). When appropriately anesthetized (i.e., no  
15 response to toe pinch), the animals' ears were marked for identification purposes. Animals were then placed on a sterile pad and their abdomen and scrotum washed with betadine and 70% alcohol. The testes were removed via a midline scrotal incision, with sterile suture being used to ligate supra-testicular tissue prior to surgical removal of each testis. The surgical wound site was closed with sterile stainless steel wound clips, and  
20 the site cleaned with betadine. The animals were allowed to recover on a sterile pad (until able to stand) and then returned to their cage.

[000109] Twenty-four hours later, animals in groups 2 through 29 were re-anesthetized with ketamine/xylazine, and an Alzet osmotic pump(s) (model 2002) was placed  
25 subcutaneously in the scapular region. In this instance, the scapular region was shaved and cleaned (betadine and alcohol) and a small incision (1 cm) made using a sterile scalpel. The osmotic pump was inserted and the wound closed with a sterile stainless steel wound clip. Animals were allowed to recover and were returned to their cage. Osmotic pumps contained the appropriate treatment (designated in Table 1) dissolved in  
30 polyethylene glycol 300 (PEG300). Osmotic pumps were filled with the appropriate solution one day prior to implantation. Animals were monitored daily for signs of acute toxicity to drug treatment (e.g., lethargy, rough coat).

[000110] After 14 days of drug treatment, rats were anesthetized with ketamine/xylazine. Animals were then sacrificed by exsanguinations under anesthesia. A blood sample was collected by venipuncture of the abdominal aorta, and submitted for complete blood cell analysis. A portion of the blood was placed in a separate tube, centrifuged at 12,000g for 1 minute, and the plasma layer removed and frozen at -20°C. The ventral prostates, seminal vesicles, levator ani muscle, liver, kidneys, spleen, lungs, and heart were removed, cleared of extraneous tissue, weighed, and placed in vials containing 10% neutral buffered formalin. Preserved tissues were sent to GTx, Inc. for histopathological analysis.

10

[000111] For data analysis, the weights of all organs were normalized to body weight, and analyzed for any statistical significant difference by single-factor ANOVA. The weights of prostate and seminal vesicle were used as indexes for evaluation of androgenic activity, and the levator ani muscle weight was used to evaluate the anabolic activity.

15

## RESULTS

[000112] The androgenic and anabolic activities the S isomers of compounds GTx-014, GTx-015, GTx-016 and GTx-007, and the R isomer of GTx-014 were examined in a castrated rat model after 14 days of administration. Testosterone propionate, at increasing doses, was used as the positive control of anabolic and androgenic effects.

20

[000113] As shown in Figures 2 and 3, the weights of prostate, seminal vesicle, and levator ani muscle in castrated, vehicle-treated rats decreased significantly, due to the ablation of endogenous androgen production. Exogenous administration of testosterone propionate, an androgenic and anabolic steroid, increased the weights of prostate, seminal vesicle, and levator ani muscle in castrated rats in a dose-dependent manner. The R-isomer of GTx-014, and S-isomers of GTx-015 and GTx-016 showed no effect on the weights of prostate, seminal vesicle, and levator ani muscle in castrated animals (data not shown). The S-isomers of GTx-007 (Figure 2: S-GTx-007) and GTx-014 (Figure 3: S-GTx-014) resulted in dose-dependent increases in prostate, seminal vesicle and levator ani muscle weights. Compared with testosterone propionate, S-GTx-007

30

showed lower potency and intrinsic activity in increasing the weights of prostate and seminal vesicle, but a greater potency and intrinsic activity in increasing the weight of levator ani muscle. Particularly, S-GTx-007, at a dose as low as 0.3 mg/day, was able to maintain the levator ani muscle weight of castrated animals in the same level as that of intact animals. Thus, S-GTx-007 is a potent nonsteroidal anabolic agent with less androgenic activity but more anabolic activity than testosterone propionate. This is a significant improvement over previous claims, in that this compound selectively stimulates muscle growth and other anabolic effects while having less effect on the prostate and seminal vesicles. This may be particularly relevant in aging men with concerns related to the development or progression of prostate cancer.

[000114] GTx-014 was less potent than GTx-007, but showed greater tissue selectivity (compare effects on the prostate and seminal vesicles in Figures 2 and 3). GTx-014 significantly increased levator ani muscle weights, but showed little to no ability to stimulate prostate and seminal vesicle growth (i.e., the prostate and seminal vesicle weights were less than 20% of that observed in intact animals or in animals treated with testosterone propionate).

[000115] Results showed that none of the examined compounds produced significant effect on body weight or the weights of other organs (i.e., liver, kidneys, spleen, lungs and heart). Nor did any compound produce any signs of acute toxicity, as gauged by diagnostic hematology tests and visual examination of animals receiving treatments. Importantly, GTx-007 did not suppress the production of luteinizing hormone (LH) or follicle stimulating hormone (FSH) at a dose of 0.3 mg/day (i.e., a dose that exhibited maximal anabolic effects).

[000116] In summary, S-GTx-007 exhibited exceptional anabolic activity in animals by maintaining the weight of levator ani muscle after removal of endogenous androgen. This discovery represents major progress towards the development of therapeutically useful nonsteroidal androgens, and a major improvement (i.e., tissue selectivity and potency) over previous drugs in this class. S-GTx-014 and S-GTx-007 showed selective anabolic activity in comparison with testosterone propionate, an androgenic and anabolic

steroid. The tissue-selective activity is actually one of the advantages of nonsteroidal androgens in terms of anabolic-related applications.

[000117] Despite similarities in structure and in-vitro functional activity, the S-isomers  
5 of compounds GTx-014, GTx-015, GTx-016, and GTx-007 exhibited profound  
differences in terms of their in-vivo activity. GTx-007 the most efficacious androgenic  
and anabolic activity in animals, with the anabolic activity greater than that of  
testosterone propionate. GTx-014 showed a small degree of androgenic activity, but an  
anabolic activity comparable to testosterone propionate. In contrast, GTx-015 and GTx-  
10 016 failed to produce any androgenic or anabolic activity in-vivo.

[000118] These studies show the discovery of two members (GTx-014 and GTx-007,  
compounds, compounds II and V respectively) of a new class of selective androgen  
receptor modulators (SARMS) that demonstrate potent anabolic effects (e.g., muscle  
15 growth) with less androgenic activity (e.g., prostatic growth). This new class of drugs  
has several advantages over non-selective androgens, including potential therapeutic  
applications in males and females for modulation of fertility, erythropoiesis,  
osteoporosis, sexual libido and in men with or at high risk for prostate cancer.

20 [000119] Further, Figures 7 and 8 demonstrate the effects of GTx-014 and GTx-007 on  
LH and FSH levels in rats. These results further demonstrate the novelty of these  
SARMS, due to their differential effects on these reproductive hormones, thus  
demonstrating the tissue-specific pharmacologic activity. In Figure 7, LH levels in  
castrated animals treated with TP and GTx-014 were significantly lower than those of  
25 untreated animals (i.e., castrated controls) at doses greater than or equal to 0.3 mg/day.  
However, higher doses (i.e., 0.5 mg/day or higher) of GTx-007 were required before  
significant decreases in LH levels were observed. Thus, GTx-007 does not suppress LH  
levels at doses that are capable of eliciting maximal stimulation of levator ani muscle  
growth. In Figure 8, FSH levels in castrated animals treated with GTx-014 were  
30 significantly lower than those of untreated animals (i.e., castrated controls) at doses of  
0.5 mg/day or higher. Similarly, lower FSH levels were observed in animals treated with  
TP. However, only this difference was only significant at a dose of 0.75 mg/day. FSH

levels in animals treated with GTx-007 were not significantly different from those of untreated animals at any dose level tested. Thus, GTx-007 does not suppress FSH levels at doses that are capable of eliciting maximal stimulation of levator ani muscle growth.

5 **Table 1. Animals Groups and Experimental Design**

Group #	Castrated?	Drug	Dose	# of animals
1	No	None	None	5
2	Yes	None	Vehicle only	5
3	Yes	Testosterone	0.1 mg/day	5
4	Yes	Testosterone	0.3 mg/day	5
5	Yes	Testosterone	0.5 mg/day	5
6	Yes	Testosterone	0.75 mg/day	5
7	Yes	Testosterone	1.0 mg/day	5
8	Yes	R-GTx-014	1.0 mg/day	5
9	Yes	S-GTx-014	0.1 mg/day	5
10	Yes	S-GTx-014	0.3 mg/day	5
11	Yes	S-GTx-014	0.5 mg/day	5
12	Yes	S-GTx-014	0.75 mg/day	5
13	Yes	S-GTx-014	1.0 mg/day	5
14	Yes	S-GTx-015	0.1 mg/day	5
15	Yes	S-GTx-015	0.3 mg/day	5
16	Yes	S-GTx-015	0.5 mg/day	5
17	Yes	S-GTx-015	0.75 mg/day	5
18	Yes	S-GTx-015	1.0 mg/day	5
19	Yes	S-GTx-016	0.1 mg/day	5
20	Yes	S-GTx-016	0.3 mg/day	5
21	Yes	S-GTx-016	0.5 mg/day	5
22	Yes	S-GTx-016	0.75 mg/day	5
23	Yes	S-GTx-016	1.0 mg/day	5
24	Yes	S-GTx-007	0.1 mg/day	5
25	Yes	S-GTx-007	0.3 mg/day	5
26	Yes	S-GTx-007	0.5 mg/day	5
27	Yes	S-GTx-007	0.75 mg/day	5
28	Yes	S-GTx-007	1.0 mg/day	5
29	Yes	None	Vehicle only	5



5

**EXAMPLE 3****Pharmacokinetics of GTx-007 in Dogs**

10 [000120] The pharmacokinetics of S-GTx-007, a novel selective androgen receptor modulator (SARM), were characterized in beagle dogs. A four-treatment, four-period crossover design was utilized in the study, which involved a total of six beagle dogs, three of each gender. Each animal received a 3 mg/kg IV dose, a 10 mg/kg IV dose, a 10 mg/kg PO dose in solution, and a 10 mg/kg PO dose in capsule, in a randomly assigned  
15 order. There was an one-week washout period between treatments. Plasma samples were collected for up to 72 hr after drug administration. Plasma S-GTx-007 concentrations were analyzed by a validated HPLC method. The clearance (CL), volume of distribution (V<sub>ss</sub>), half-life (T<sub>1/2</sub>), and other pharmacokinetic parameters were determined by noncompartmental methods. Results showed that S-GTx-007 was cleared from dog  
20 plasma with a terminal T<sub>1/2</sub> of about 4 hr and a CL of 4.4 mL/min/kg after IV administration. Figures 4, 5, and 6 show the plasma concentration-time profiles of S-GTx-007 after administration of an intravenous solution, oral solution, and oral capsule, respectively. The pharmacokinetics were dose- and gender-independent. The oral bioavailability of S-GTx-007 varied with the dosage form, and averaged 38% and 19%  
25 for solution and capsule, respectively. Thus, S-GTx-007 demonstrated moderate half-life, slow clearance and moderate bioavailability in beagle dogs, identifying it as the first of a new class of orally bioavailable tissue-selective androgen receptor modulators.

**EXAMPLE 4****GTx-007 Analysis by HPLC**

[000121] A reversed phase high pressure liquid chromatograph (HPLC) assay was


developed to quantitate GTx-007 concentrations in dog plasma. Dog blood samples were obtained by venipuncture and centrifuged at 1000g for 15 minutes. Samples were stored frozen at -20°C until analysis. Individual samples were rapidly thawed and an aliquot (0.5 ml) was spiked with internal standard (20 µl of a 200 µg/ml aqueous solution of CM-II-87). An aliquot of 1 ml of acetonitrile was added to the samples to precipitate plasma proteins. The samples were vortexed and then centrifuged at 1000g for 15 minutes. The supernatant was decanted into glass extraction tubes and 7.5 ml of ethyl acetate was added. The extraction mixture was left at room temperature for 20 minutes, and vortexed several times during this interval. The samples were then centrifuged at 1000g for 10 minutes, and the organic phase was removed and placed in conical-bottomed glass tubes. The organic phase was evaporated under nitrogen. The samples were reconstituted in 200 µl of mobile phase (35:65 acetonitrile:water) and transferred to an autosampler vial for HPLC injection (Waters 717 plus autosampler, Waters Corp., Milford, MA). The isocratic mobile phase of 35% (v/v) acetonitrile in water was pumped at a flow rate of 1 ml/min (Model 510, Waters Corp.). The stationary phase was a C18 reversed phase column (Novapak C18, 3.9 x 150 mm). Analytes were monitored with UV detection at 270 nm (Model 486 absorbance detector, Waters Corp.). Retention times for GTx-007 and CM-II-87 were 11.1 and 16.9 minutes, respectively. Chromatography data was collected and analyzed using Millennium software. Plasma concentrations of GTx-007 in each sample were determined by comparison to calibration curves. Calibration curves were constructed by adding known amounts of GTx-007 to dog plasma. Final GTx-007 concentrations in dog plasma samples used in the calibration curves were 0.08, 0.2, 0.4, 2, 4, 10, and 20 µg/ml. Calibration curves were linear over this concentration range and exhibited correlation coefficients ( $r^2$ ) of 0.9935 or greater. Intra- and inter-day coefficients of variation for the standards ranged from 6.4% for 0.08 µg/ml to 7.9% for 20 µg/ml.

[000122] Melting points were determined on a Thomas-Hoover capillary melting point apparatus and are uncorrected. Infrared spectra were recorded on a Perkin Elmer System 2000 FT-IR. Optical rotations were determined on an Autopol<sup>®</sup> III Automatic Polarimeter (Rudolph Research Model III-589-10, Fairfield, New Jersey). Proton and carbon-13 magnetic resonance spectra were obtained on a Bruker AX 300 spectrometer

(300 and 75 MHz for  $^1\text{H}$  and  $^{13}\text{C}$ , respectively). Chemical shift values were reported as parts per million ( $\delta$ ) relative to tetramethylsilane (TMS). Spectral data were consistent with assigned structures. Mass spectra were determined on a Bruker-HP Esquire LC System. Elemental analyses were performed by Atlantic Microlab Inc. (Norcross, GA),  
5 and found values were within 0.4 % of the theoretical values. Routine thin-layer chromatography (TLC) was performed on silica gel on aluminum plates (silica gel 60 F 254, 20 x 20 cm, Aldrich Chemical Company Inc., Milwaukee, WI). Flash chromatography was performed on silica gel (Merck, grade 60, 230-400 mesh, 60). Tetrahydrofuran (THF) was dried by distillation over sodium metal. Acetonitrile  
10 (MeCN) and methylene chloride ( $\text{CH}_2\text{Cl}_2$ ) were dried by distillation from  $\text{P}_2\text{O}_5$ .

WHAT IS CLAIMED IS:

1. A selective androgen receptor modulator compound having tissue-selective in-  
 5 vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor  
 represented by the structure of formula I:




wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;


Z is NO<sub>2</sub>, CN, COR, COOH or CONHR;

10 Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;


Q is alkyl, halogen, NR<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR,  
 NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR, NHSO<sub>2</sub>CH<sub>3</sub>,  
 NHSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR wherein R is an alkyl, aryl,  
 C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl; or Q together  
 15 with the benzene ring to which it is attached is a fused ring system represented  
 by structure A, B or C:



A



B



C

$R_1$  is  $\text{CH}_3$ ,  $\text{CF}_3$ ,  $\text{CH}_2\text{CH}_3$ , or  $\text{CF}_2\text{CF}_3$ ; and

T is OH, OR,  $-\text{NHCOCH}_3$ , or  $\text{NHCOR}$  wherein R is a  $\text{C}_1$ - $\text{C}_4$  alkyl, a  $\text{C}_1$ - $\text{C}_4$  haloalkyl, phenyl, halo, alkenyl or hydroxyl.

2. The selective androgen receptor modulator compound of claim 1, wherein Q is in the para position.

3. The selective androgen receptor modulator compound, of claim 2, wherein Q is alkyl, halogen,  $\text{NR}_2$ ,  $\text{NHCOCH}_3$ ,  $\text{NHCOCF}_3$ ,  $\text{NHCOR}$ ,  $\text{NHCONHR}$ ,  $\text{NHCOOR}$ ,  $\text{OCONHR}$ ,  $\text{CONHR}$ ,  $\text{NHCSCH}_3$ ,  $\text{NHCSCF}_3$ ,  $\text{NHCSR N HSO}_2\text{CH}_3$ ,  $\text{NHSO}_2\text{R}$ , OR, COR, OCOR,  $\text{OSO}_2\text{R}$ ,  $\text{SO}_2\text{R}$  or SR wherein R is an alkyl, aryl,  $\text{C}_1$ - $\text{C}_4$  alkyl, a  $\text{C}_1$ - $\text{C}_4$  haloalkyl, phenyl, halo, alkenyl or hydroxyl.

4. The selective androgen receptor modulator compound of claim 1, wherein X is O.

5. The selective androgen receptor modulator compound of claim 4, wherein Q is in the para position.

6. The selective androgen receptor modulator compound, of claim 5, wherein Q is alkyl, halogen,  $\text{NR}_2$ ,  $\text{NHCOCH}_3$ ,  $\text{NHCOCF}_3$ ,  $\text{NHCOR}$ ,  $\text{NHCONHR}$ ,  $\text{NHCOOR}$ ,  $\text{OCONHR}$ ,  $\text{CONHR}$ ,  $\text{NHCSCH}_3$ ,  $\text{NHCSCF}_3$ ,  $\text{NHCSR N HSO}_2\text{CH}_3$ ,  $\text{NHSO}_2\text{R}$ , OR, COR, OCOR,  $\text{OSO}_2\text{R}$ ,  $\text{SO}_2\text{R}$  or SR wherein R is a  $\text{C}_1$ - $\text{C}_4$  alkyl, a  $\text{C}_1$ - $\text{C}_4$  haloalkyl, phenyl, halo, alkenyl or hydroxyl.

7. The selective androgen receptor modulator compound of claim 1, wherein Z is para NO<sub>2</sub>.

8. The selective androgen receptor modulator compound of claim 1, wherein Y is meta CF<sub>3</sub>.


9. The selective androgen receptor modulator compound of claim 1, wherein T is OH.

10. The selective androgen receptor modulator compound of claim 1, wherein R<sub>1</sub> is CH<sub>3</sub>.

11. The selective androgen receptor modulator compound of claim 1, wherein Q is para F.

15


12. The selective androgen receptor modulator compound of claim 1, represented by the structure of formula IV.



IV

20 13. The selective androgen receptor modulator compound of claim 1, wherein Q is para COCH<sub>3</sub>.


14. The selective androgen receptor modulator compound of claim 1, represented by the structure of formula V.



V 55

15. The selective androgen receptor modulator compound of claim 1, wherein Q is para COCH<sub>2</sub>CH<sub>3</sub>.


- 5 16. The selective androgen receptor modulator compound of claim 1, represented by the structure of formula VI.



VI

17. The selective androgen receptor modulator compound of claim 1, wherein Q is para NHCOCH<sub>3</sub>.

- 10 18. The selective androgen receptor modulator compound of claim 1, represented by the structure of formula VII.



VII

- 15 19. A composition comprising the selective androgen receptor modulator compound of claim 1 and a suitable carrier or diluent alone or in combination with a progestin or estrogen.

- 20 20. A pharmaceutical composition comprising an effective amount of the selective androgen receptor modulator compound of claim 1 alone or in combination with a progestin or estrogen and a pharmaceutically acceptable carrier, diluent or salt.

21. A method of binding a selective androgen receptor modulator compound to a androgen receptor comprising contacting the androgen receptor with the selective androgen receptor modulator compound of claim 1 under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor.

5

22. A method of modulating spermatogenesis in a subject comprising contacting an androgen receptor of the subject with the selective androgen receptor modulator compound of claim 1 under conditions effective to increase or decrease sperm production.

10

23. A method of hormone therapy comprising contacting an androgen receptor of a subject with the selective androgen receptor modulator compound of claim 1 under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor and effect a change in an androgen-dependent condition.

15


24. A method of treating a subject having a hormone related condition comprising contacting an androgen receptor of a subject with the selective androgen receptor modulator compound of claim 1 under conditions effective to bind the selective androgen receptor modulator compound to the androgen receptor and effect a change in an androgen-dependent condition.

20

25. A method of treating a subject having prostate cancer comprising administering to said subject an effective amount of a selective androgen receptor modulator compound according to claim 1.

25

26. A selective androgen receptor modulator compound having in-vivo androgenic and anabolic activity of a nonsteroidal ligand for the androgen receptor having the formula III:






where X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;  
Z is NO<sub>2</sub>, CN, COR, or CONHR;  
Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>;  
5 R is an alkyl group or OH; and  
Q is acetamido or trifluoroacetamido.

27. The selective androgen receptor modulator compound of claim 26, wherein X is O.
- 10 28. The selective androgen receptor modulator compound of claim 26, wherein Z is NO<sub>2</sub>.
29. The selective androgen receptor modulator compound of claim 26, wherein Y is CF<sub>3</sub>.
- 15 30. The selective androgen receptor modulator compound of claim 26, wherein Q is NHCOCH<sub>3</sub>.
31. The selective androgen receptor modulator compound of claim 26, wherein X is O, Z is NO<sub>2</sub>, Y is CF<sub>3</sub>, and Q is NHCOCH<sub>3</sub>.
32. A composition comprising the selective androgen receptor modulator compound of claim 26 and a suitable carrier or diluent.
- 20 33. A pharmaceutical composition comprising an effective amount of the selective androgen receptor modulator compound of claim 26 and a pharmaceutically acceptable carrier, diluent or salt.
34. A method of binding a selective androgen receptor modulator compound to a androgen receptor comprising contacting the androgen receptor with the selective androgen receptor modulator compound of claim 26 effective to bind the selective androgen receptor modulator compound to the androgen receptor.
- 25 35. A method of suppressing spermatogenesis in a subject comprising contacting an androgen receptor of the subject with the selective androgen receptor modulator compound of claim 26 effective to suppress sperm production.
- 30 36. A method of hormone therapy comprising contacting an androgen receptor of a subject with the selective androgen receptor modulator compound of claim 26 effective

to bind the selective androgen receptor modulator compound to the androgen receptor and effect a change in an androgen-dependent condition.

37. A method of treating a subject having a hormone related condition comprising contacting an androgen receptor of a subject with the selective androgen receptor modulator compound of claim 26 effective to bind the selective androgen receptor  
5 modulator compound to the androgen receptor and effect a change in an androgen-dependent condition.

38. A nonsteroidal agonist compound represented by the structure of formula VIII:



10


wherein X is a O, CH<sub>2</sub>, NH, Se, PR, or NR;

R<sub>1</sub> is CH<sub>3</sub>, CF<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, or CF<sub>2</sub>CF<sub>3</sub>;


T is OH, OR, -NHCOCH<sub>3</sub>, or NHCOR wherein R is a C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl;

15

A is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or heterocyclic ring represented by the structure:



B is a 5 or 6 membered saturated, unsaturated or aromatic carbocyclic or  
20 heterocyclic ring represented by the structure:



wherein

$A_1 - A_{11}$  are each C, O, S or N;

$B_1 - B_{11}$  are each C, O, S or N;

Z is NO<sub>2</sub>, CN, COR, COOH or CONHR;

Y is I, CF<sub>3</sub>, Br, Cl, or SnR<sub>3</sub>; and

$Q_1$  and  $Q_2$  are independently of each other alkyl, halogen,  $NR_2$ ,

NHCOCH<sub>3</sub>, NHCOCF<sub>3</sub>, NHCOR, NHCONHR, NHCOOR, OCONHR, CONHR, NHCSCH<sub>3</sub>, NHCSCF<sub>3</sub>, NHCSR NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>R, OR, COR, OCOR, OSO<sub>2</sub>R, SO<sub>2</sub>R or SR wherein R is a C<sub>1</sub>-C<sub>4</sub> alkyl, a C<sub>1</sub>-C<sub>4</sub> haloalkyl, phenyl, halo, alkenyl or hydroxyl.

39. A composition comprising the non-steroidal agonist compound of claim 38 and a suitable carrier or diluent alone or in combination with a progestin or estrogen.

40. A pharmaceutical composition comprising an effective amount of the non-steroidal agonist compound of claim 38 alone or in combination with a progestin or estrogen and a pharmaceutically acceptable carrier, diluent or salt.

41. A method of binding a compound to an androgen receptor comprising contacting the androgen receptor with the non-steroidal agonist compound of claim 38 under conditions effective to bind the non-steroidal agonist compound to the androgen receptor.

42. A method of modulating spermatogenesis in a subject comprising contacting an androgen receptor of the subject with the non-steroidal agonist compound of claim 38 under conditions effective to increase or decrease sperm production.

43. A method of hormone therapy comprising contacting an androgen receptor of a subject with the non-steroidal agonist compound of claim 38 under conditions effective

to bind the non-steroidal agonist compound to the androgen receptor and effect a change in an androgen-dependent condition.

44. A method of treating a subject having a hormone related condition comprising  
5 contacting an androgen receptor of a subject with the non-steroidal agonist compound of claim 38 under conditions effective to bind the non-steroidal agonist compound to the androgen receptor and effect a change in an androgen-dependent condition.

45. A method of treating a subject having prostate cancer comprising administering  
10 to said subject an effective amount of the non-steroidal agonist compound according to claim 38.

46. A method of determining the presence of the compound of claim 1 in a sample, said method comprising the steps of:  
15 obtaining said sample; and  
detecting said compound in said sample;  
thereby determining the presence of said compound in said sample.

47. The method according to claim 46, wherein the sample is a blood serum,  
20 plasma, urine, or saliva sample.

48. The method according to claim 46, wherein the detection step comprises measuring the absorbance of the compound.

25 49. A method of determining the presence of the compound of claim 26 in a sample, said method comprising the steps of:  
obtaining said sample; and  
detecting said compound in said sample;  
thereby determining the presence of said compound in said sample.

30 50. The method according to claim 49, wherein the sample is a blood serum, plasma, urine, or saliva sample.

51. The method according to claim 49, wherein the detection step comprises measuring the absorbance of the compound.

5 52. A method of determining the presence of the compound of claim 38 in a sample, said method comprising the steps of:

obtaining said sample; and

detecting said compound in said sample;

thereby determining the presence of said compound in said sample.

10

53. The method according to claim 52, wherein the sample is a blood serum, plasma, urine, or saliva sample.

15 54. The method according to claim 52, wherein the detection step comprises measuring the absorbance of the compound.

1/9




FIG.1

2/9




FIG.2




FIG. 3



4/9




FIG. 4

5/9




FIG.5

6/9




FIG.6

7/9




FIG. 7

8/9




FIG.8

9/9






FIG. 9

## INTERNATIONAL SEARCH REPORT

 International application No.  
 PCT/US01/26328

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(7) : Please See Extra Sheet. US CL : 514/493, 616, 619; 564/153, 156, 157, 158 According to International Patent Classification (IPC) or to both national classification and IPC																				
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/493, 616, 619; 564/153, 156, 157, 158 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched NONE Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAPLUS on STN																				
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>																				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.																		
Y	US 4,636,505 A (TUCKER) 13 January 1987 (13/01/87), see entire document.	1-54																		
A	EP 0 253 503 A2 (IMPERIAL CHEMICAL INDUSTRIES PLC) 20 January 1988 (20/01/88), see entire document.	1-54																		
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.																				
<table border="0"> <tr> <td colspan="2">* Special categories of cited documents</td> <td>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"A" document defining the general state of the art which is not considered to be of particular relevance</td> <td>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> <td></td> </tr> <tr> <td>"E" earlier document published on or after the international filing date</td> <td>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> <td></td> </tr> <tr> <td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"A" document member of the same patent family</td> <td></td> </tr> <tr> <td>"O" document referring to an oral disclosure, use, exhibition or other means</td> <td></td> <td></td> </tr> <tr> <td>"P" document published prior to the international filing date but later than the priority date claimed</td> <td></td> <td></td> </tr> </table>			* Special categories of cited documents		"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone		"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art		"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A" document member of the same patent family		"O" document referring to an oral disclosure, use, exhibition or other means			"P" document published prior to the international filing date but later than the priority date claimed		
* Special categories of cited documents		"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention																		
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone																			
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art																			
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A" document member of the same patent family																			
"O" document referring to an oral disclosure, use, exhibition or other means																				
"P" document published prior to the international filing date but later than the priority date claimed																				
Date of the actual completion of the international search 20 SEPTEMBER 2001		Date of mailing of the international search report 16 NOV 2001																		
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer BRIAN J. DAVIS <i>Brian J. Davis</i> Telephone No. (703) 308-1235																		

INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US01/26328

A. CLASSIFICATION OF SUBJECT MATTER:  
IPC (7):

C07C 233/15, 233/33, 233/48, 235/26, 237/14; A01N 37/18; A01K 31/16, 31/165